

Entirety of cuspidal Eisenstein series on loop groups

Howard Garland, Stephen D. Miller*, and Manish M. Patnaik†

April 17, 2013

Abstract

In this paper, we prove the entirety of loop group Eisenstein series induced from cusp forms on the underlying finite dimensional group, by demonstrating their absolute convergence on the full complex plane. This is quite in contrast to the finite-dimensional setting, where such series only converge absolutely in a right half plane (and have poles elsewhere coming from L -functions in their constant terms). Our result is the $\mathbb{Q}$ -analog of a theorem of A. Braverman and D. Kazhdan from the function field setting [4, Theorem 5.2], who previously showed the analogous Eisenstein series are finite sums.

1 Introduction

The theory of Eisenstein series on finite dimensional Lie groups has had profound applications in number theory, geometry, and mathematical physics. To name one important example, Langlands' monograph *Euler Products* [24] gave meromorphic continuations to the new L -functions he discovered in their constant terms. This work not only led him to formulate his functoriality conjectures, but also initiated the Langlands-Shahidi method of constructing and analyzing L -functions. The method frequently obtains the most powerful results on analytic continuations of L -functions we presently know, but does not treat all L -functions; instead, it only applies when an underlying representation occurs in the adjoint action of the Levi component of a parabolic on its nilradical. Some of the most fascinating and important applications to number theory and the Langlands program come from maximal parabolic subgroups of exceptional groups. Unfortunately, these are finite in number and all L -functions under the purview of the Langlands-Shahidi method have already been treated.

It has been an intriguing possibility to expand the landscape of possible parabolic subgroups by considering infinite-dimensional Kac-Moody groups such as loop groups. This setting features a vastly wider range of Levi adjoint actions (some of which we have identified and which will be the subject of a future paper), and thus the possibility of new L -functions. This paper is a continuation of a series of papers [8–14] by the first-named author on the loop Eisenstein series.

Loop Eisenstein series induced from quasi-characters on a minimal parabolic subgroup were shown to converge in a region analogous to the shifted Weyl chamber established by Godement [15, §3] in the finite-dimensional setting. By developing an analog of the Maass-Selberg relations,

*Supported by NSF grant DMS-1201362

†Supported in part by an NSF Postdoctoral Fellowship DMS-0802940 and an University of Alberta Startup grant

they were then meromorphically continued to a larger region in [10–13]. In contrast to the finite dimensional case, however, these minimal parabolic loop Eisenstein series cannot be meromorphically continued everywhere, and it was observed in [8] that there is a natural boundary or “wall” beyond which the series cannot be continued.

Because the Levi components of parabolic subgroups of loop groups are finite dimensional, there are also *cuspidal loop Eisenstein series* induced from cusp forms on these classical Levi components. In [14], their convergence in the Godement-type region was deduced by a domination argument from the convergence of the minimal-parabolic loop Eisenstein series. Moreover, their extension up to the location of the aforementioned “wall” was also shown using the Maass-Selberg relations [10–13]. Somewhat surprisingly, as was observed in [14], the Maass-Selberg relations here consist of just a single holomorphic term, and consequently these cuspidal Eisenstein series have no poles in this region.

In this paper we show a much stronger result holds in the case of cuspidal loop Eisenstein series, at least those induced from the maximal parabolic whose Levi component comes from the underlying finite-dimensional root system. The following result analytically continues these series beyond the “wall”:

Theorem. *Let ϕ be a spherical cusp form for a Chevalley group G over $\mathbb{Q}$. Then the cuspidally-induced Eisenstein series $E_{\phi, \nu}$ (4.27) on the loop group of G converges absolutely for any $\nu \in \mathbb{C}$ to an entire function of ν .*

For example, this theorem applies to Eisenstein series on the loop group E_9 induced from cusp forms ϕ on E_8 . By *spherical* we mean a cusp form which is K -fixed at all places of $\mathbb{Q}$. This restriction is undoubtedly not essential to our proof, but is made because of an obstacle coming from the local theory of loop groups: the usual definition of K -finite Eisenstein series (e.g., [31, (6.3.1)]) relies on a matrix coefficient construction whose generalization to loop groups has not been developed. Thus the loop Eisenstein series induced from K -finite cusp forms have not even been defined, and so it is premature to study their convergence. The K -finite condition at nonarchimedean places amounts to a congruence group condition, and hence a smaller group of summation in the Eisenstein series definition. Since our arguments involve dominating by larger, convergent sums, relaxing the K -fixed condition is therefore likely to make the convergence only easier.

The K -fixed condition at the archimedean place also arises in a paper by Krötz and Opdam [23] on Bernstein’s theorem that cusp forms decay exponentially. Though those methods apply to K -finite cusp forms as well (e.g., in unpublished notes by those authors), there is no proof in the literature. These decay results imply our important ingredient Theorem 4.6. For completeness, and in order to allow our convergence argument to apply to K -finite Eisenstein series once their definition has been given, we have included a proof of Theorem 4.6 for arbitrary K -finite forms in the Appendix. Together, this provides the full analytic machinery anticipated to be necessary to handle the convergence of the presently-undefined K -finite Eisenstein series. The appendix also discusses some related questions about decay estimates posed by string theorists.

A function-field analog of this result was announced by A. Braverman and D. Kazhdan in [4, Theorem 5.2(3)]. Their proof relies on a geometric interpretation of the Eisenstein series unavailable over number fields. In fact, at each fixed element in the appropriate symmetric space, their cuspidal Eisenstein series is a finite sum (independent of the spectral parameter). This finiteness crucially uses the classical theorem of G. Harder that cusp forms vanish outside of a compact subset of the fundamental domain. This vanishing outside of a compact set is far from true in the

number field case, where such forms instead merely decay rapidly in the cusps. While rapid (e.g., polynomial) decay statements are well-known, it was somewhat striking to us that the classical decay results were insufficient for our purposes (see Remark 4.6.1 for further details). Instead we require the *exponential decay* result Theorem A.1 (or its equivalent statement Theorem 4.6) mentioned above to obtain convergence in the number field setting.

Another key ingredient in our work is Theorem 3.3 and its Corollary 3.4, a growth estimate on the diagonal Iwasawa components in the classical and central directions as one varies over a discrete group. In the function field setting, this estimate – coupled with Harder’s compact support theorem – is enough to deduce the Braverman-Kazhdan result. In the number field setting, the analysis is more involved and the entirety result we prove here is instead shown by establishing a reduction to the half-plane convergence result [14] recalled in Theorem 4.23.

Acknowledgements: We would like to thank Alexander Braverman for informing us of his results with David Kazhdan and of Bernstein’s exponential decay estimates. We would also like to thank William Casselman, Michael B. Green, Bernhard Krötz, and Wilfried Schmid for their helpful comments about rapid decay.

2 Notation

In this section we shall recall some background material which is used later in the paper. Part 2A concerns the finite-dimensional situation, part 2B discusses affine loop groups, part 2C focuses on the Weyl group, and part 2D summarizes some useful decompositions. Finally, part 2E redescribes some of this material from the adelic point of view. A general reference for most of this material is Kac’s book [22].

A. Finite Dimensional Lie Algebras and Groups

2.1. Let $\mathbf{C}$ be an irreducible, classical, Cartan matrix of size $\ell \times \ell$. Let $\mathfrak{g}$ be the corresponding real, split, simple, finite-dimensional Lie algebra of rank ℓ . Choose a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$ and denote the set of roots with respect to $\mathfrak{h}$ by Δ . Let $\Pi = \{\alpha_1, \dots, \alpha_\ell\} \subset \mathfrak{h}^*$ denote a fixed choice of positive simple roots, and $\Pi^\vee = \{h_1, \dots, h_\ell\} \subset \mathfrak{h}$ the corresponding set of simple coroots. We let Δ_+ and Δ_- be the sets of positive and negative roots, respectively. For a root $\alpha \in \Delta$ we denote by h_α the corresponding coroot. Let $(\cdot, \cdot)$ denote the Killing form on $\mathfrak{h}$ normalized so that $(\alpha_0, \alpha_0) = 2$, where α_0 is the highest root of $\mathfrak{g}$, and write $\langle \cdot, \cdot \rangle$ for the natural pairing between $\mathfrak{h}^*$ and $\mathfrak{h}$.

Let W be the corresponding Weyl group of the root system defined above and Q and $Q^\vee$ the root and coroot lattices, which are spanned over $\mathbb{Z}$ by Π and $\Pi^\vee$, respectively. We define $\Lambda \supset Q$ and $\Lambda^\vee \supset Q^\vee$ to be the weight and coweight lattices, respectively. The fundamental weights will be denoted by ω_i for $i = 1, \dots, \ell$; recall that these are defined by the conditions that

$$\langle \omega_i, h_j \rangle = \begin{cases} 1, & i = j \\ 0, & i \neq j \end{cases} \quad \text{for } 1 \leq i, j \leq \ell. \quad (2.1)$$

We also define the fundamental coweights $\omega_j^\vee \in \mathfrak{h}$ for $j = 1, \dots, \ell$ by the conditions that

$$\langle \alpha_i, \omega_j^\vee \rangle = \begin{cases} 1, & i = j \\ 0, & i \neq j \end{cases} \quad \text{for } 1 \leq i, j \leq \ell. \quad (2.2)$$

Note that we have

$$(\omega_i^\vee, h_j) = \begin{cases} \frac{2}{(\alpha_i, \alpha_i)}, & i = j \\ 0, & i \neq j \end{cases} \quad \text{for } 1 \leq i, j \leq \ell. \quad (2.3)$$

As usual we set

$$\rho = \frac{1}{2} \sum_{\alpha \in \Delta_+} \alpha = \sum_{j=1}^{\ell} \omega_j, \quad (2.4)$$

which satisfies the condition

$$\langle \rho, h_i \rangle = 1 \text{ for } 1 \leq i \leq \ell. \quad (2.5)$$

We record here the following elementary statement which will be useful later. It is equivalent to the positivity of the inverse of the Cartan matrix.

Lemma ([30, p.111, Lemma 6]). *Each fundamental weight $\omega \in \{\omega_1, \dots, \omega_\ell\}$ can be written as a strictly positive linear combination of positive simple roots $\alpha_1, \dots, \alpha_\ell$.*

2.2. Let G be a finite dimensional, simple, real Chevalley group with Lie algebra $\mathfrak{g}$. Let $B \subset G$ be a Borel subgroup with unipotent radical U and split torus T . Let $A \subset T$ be the connected component of the identity of T , and assume (as we may) that B is arranged so that $\text{Lie}(A) = \mathfrak{h}$ and that $\text{Lie}(U)$ is spanned by the root vectors for Δ_+ . Then G has a maximal compact subgroup K such that the Iwasawa decomposition

$$G = UAK \quad (2.6)$$

holds with uniqueness of expression. We denote the natural projection onto the A -factor by the map $g \mapsto \text{Iw}_A(g)$. Any linear functional $\lambda : \mathfrak{h} \rightarrow \mathbb{C}$ gives rise to a quasi-character $a \mapsto a^\lambda$ of A by the formula

$$a^\lambda := \exp(\langle \lambda, \ln a \rangle). \quad (2.7)$$

2.3. Let $\Gamma \subset G$ denote the stabilizer of the fixed Chevalley lattice defining G . We then use the following notion of Siegel set as defined in [2]. Let $U_{\mathcal{D}} \subset U$ denote a fundamental domain for the action of $\Gamma \cap U$ on U . For any $t > 0$ we set

$$A_t := \{a \in A \mid a^{\alpha_i} > t \text{ for each } 1 \leq i \leq \ell\}. \quad (2.8)$$

A Siegel set has the form $\mathfrak{S}_t = U_{\mathcal{D}} A_t K$. The following is shown in [2].

Proposition. *Suppose that $t < \sqrt{3}/2$. Then for every $g \in G$ there exists $\gamma \in \Gamma$ such that $\gamma g \in \mathfrak{S}_t$, i.e., the Γ -translates of $\mathfrak{S}_t$ cover G .*

2.3.1. Remark One of the key points in the proof of this proposition is the following minimum principle (which will be important for us later). Let V^ρ be the highest weight representation of G corresponding to the dominant integral weight $\rho : \mathfrak{h} \rightarrow \mathbb{C}$ from (2.4). This representation comes equipped with a K -invariant norm $\|\cdot\|$. Let $v_\rho \in V^\rho$ denote a highest weight vector. Then for any $g \in G$, it is not hard to see that the function

$$\begin{aligned} \Psi_g : \Gamma &\rightarrow \mathbb{R}_{>0}, \\ \gamma &\mapsto \|\gamma g v_\rho\| \end{aligned} \quad (2.9)$$

achieves a minimum value (see [2]). Furthermore, if $\gamma \in \Gamma$ minimizes $\Psi_g(\gamma)$, then $\gamma g \in \mathfrak{S}_t$.

B. Affine Lie Algebras

2.4. Let $\widehat{\mathbf{C}}$ denote the $(\ell + 1) \times (\ell + 1)$ affine Cartan matrix corresponding to $\mathbf{C}$, and $\widehat{\mathfrak{g}}$ the one-dimensional central extension of the loop algebra corresponding to the finite dimensional Lie algebra $\mathfrak{g}$ associated to $\mathbf{C}$. The extension

$$\widehat{\mathfrak{g}}^e = \widehat{\mathfrak{g}} \oplus \mathbb{R}\mathbf{D} \quad (2.10)$$

of $\widehat{\mathfrak{g}}$ by the *degree derivation* $\mathbf{D}$ is the (untwisted) affine Kac-Moody algebra associated to $\widehat{\mathbf{C}}$ (see [Kac, Chapter 6, 7]).

Let $\widehat{\mathfrak{h}} \subset \widehat{\mathfrak{g}}$ be a Cartan subalgebra and define the extended Cartan subalgebra

$$\widehat{\mathfrak{h}}^e := \widehat{\mathfrak{h}} \oplus \mathbb{R}\mathbf{D}. \quad (2.11)$$

Let $(\widehat{\mathfrak{h}}^e)^*$ denote the (real) algebraic dual of $\widehat{\mathfrak{h}}^e$ and continue to denote the natural pairing as

$$\langle \cdot, \cdot \rangle : (\widehat{\mathfrak{h}}^e)^* \times \widehat{\mathfrak{h}}^e \rightarrow \mathbb{R}. \quad (2.12)$$

The set of simple affine roots will be denoted by

$$\widehat{\Pi} = \{a_1, \dots, a_{\ell+1}\} \subset (\widehat{\mathfrak{h}}^e)^*. \quad (2.13)$$

Similarly, we write

$$\widehat{\Pi}^\vee = \{h_1, \dots, h_{\ell+1}\} \subset \widehat{\mathfrak{h}} \subset \widehat{\mathfrak{h}}^e \quad (2.14)$$

for the set of simple affine coroots. Note that the simple roots $a_i : \widehat{\mathfrak{h}}^e \rightarrow \mathbb{R}$ satisfy the relations

$$\begin{aligned} \langle a_i, \mathbf{D} \rangle &= 0 \quad \text{for } i = 1, \dots, \ell \\ \text{and } \langle a_{\ell+1}, \mathbf{D} \rangle &= 1. \end{aligned} \quad (2.15)$$

A more explicit description of the a_i will be given in section 2.5.

We define the affine root lattice as $\widehat{Q} = \mathbb{Z}a_1 + \dots + \mathbb{Z}a_{\ell+1}$ and the affine coroot lattice as $\widehat{Q}^\vee = \mathbb{Z}h_1 + \dots + \mathbb{Z}h_{\ell+1}$. We shall denote the subset of non-negative integral linear combinations of the a_i (respectively, h_i) as $\widehat{Q}_+$ (respectively, $\widehat{Q}_+^\vee$). The integral weight lattice is defined as

$$\widehat{\Lambda} := \{\lambda \in \widehat{\mathfrak{h}}^* \mid \langle \lambda, h_i \rangle \in \mathbb{Z} \text{ for } i = 1, \dots, \ell+1\}. \quad (2.16)$$

We regard $\widehat{\Lambda}$ as a subset of $(\widehat{\mathfrak{h}}^e)^*$ by declaring that

$$\langle \lambda, \mathbf{D} \rangle = 0 \quad \text{for } \lambda \in \widehat{\Lambda}. \quad (2.17)$$

The lattice $\widehat{\Lambda}$ is spanned by the fundamental affine weights $\Lambda_1, \dots, \Lambda_{\ell+1}$, which are defined by the conditions that

$$\langle \Lambda_i, h_j \rangle = \begin{cases} 1, & i = j \\ 0, & i \neq j \end{cases} \quad \text{for } 1 \leq i, j \leq \ell+1. \quad (2.18)$$

Note that the dual space $(\widehat{\mathfrak{h}}^e)^*$ is spanned by $a_1, a_2, \dots, a_{\ell+1}, \Lambda_{\ell+1}$.

For any $a \in \widehat{Q}$ define

$$\widehat{\mathfrak{g}}^a := \{x \in \widehat{\mathfrak{g}} \mid [h, x] = \langle a, h \rangle x, \text{ for all } h \in \widehat{\mathfrak{h}}\}. \quad (2.19)$$

The set of nonzero $a \in \widehat{Q}$ such that $\dim \widehat{\mathfrak{g}}^a > 0$ will be called the roots of $\widehat{\mathfrak{g}}$ and denoted by $\widehat{\Delta}$. The sets $\widehat{\Delta}_+ := \widehat{\Delta} \cap \widehat{Q}_+$ and $\widehat{\Delta}_- := \widehat{\Delta} \cap (-\widehat{Q}_+)$ will be called the sets of positive and negative affine roots, respectively. We have that $\widehat{\Delta} = \widehat{\Delta}_+ \sqcup \widehat{\Delta}_-$, i.e., every element in $\widehat{\Delta}$ can be written as a linear combination of elements from $\widehat{\Pi}$ with all positive integral or all negative integral coefficients.

For each $i \in \{1, 2, \dots, \ell + 1\}$ we define the reflection $w_i : (\widehat{\mathfrak{h}}^e)^* \rightarrow (\widehat{\mathfrak{h}}^e)^*$ by the formula

$$w_i : \lambda \mapsto \lambda - \langle \lambda, h_i \rangle a_i \quad \text{for } \lambda \in (\widehat{\mathfrak{h}}^e)^*. \quad (2.20)$$

The Weyl group $\widehat{W} \subset \text{Aut}((\widehat{\mathfrak{h}}^e)^*)$ is the group generated by the elements $w_1, \dots, w_{\ell+1}$. The dual action of $\widehat{W}$ on $\widehat{\mathfrak{h}}^e$ is defined by the formula

$$\langle \lambda, w \cdot h \rangle = \langle w^{-1} \lambda, h \rangle \quad \text{for all } \lambda \in (\widehat{\mathfrak{h}}^e)^*, h \in \widehat{\mathfrak{h}}^e, \text{ and } w \in \widehat{W}. \quad (2.21)$$

The roots decompose as

$$\widehat{\Delta} = \widehat{\Delta}_W \sqcup \widehat{\Delta}_I, \quad (2.22)$$

where $\widehat{\Delta}_W$ (known as the “real roots” or “Weyl roots”) is the $\widehat{W}$ -orbit of $\widehat{\Pi}$, and $\widehat{\Delta}_I$ (known as the “imaginary roots”) is its complement in $\widehat{\Delta}$. These sets will be described explicitly in (2.29)-(2.30) below. Each imaginary root is fixed by $\widehat{W}$. The space $\widehat{\mathfrak{g}}^a$ is 1-dimensional for $a \in \widehat{\Delta}_W$ and is ℓ -dimensional for $a \in \widehat{\Delta}_I$. Coroots of elements $a \in \widehat{\Delta}_W$ can be defined by the formula

$$h_a := w^{-1} h_j \in \widehat{\mathfrak{h}}, \quad (2.23)$$

where w is an element of $\widehat{W}$ such that $wa = a_j$ (this is shown to be well-defined in [22, §5.1]).

2.5. We shall now give a more concrete description of the affine roots and coroots in terms of the underlying finite dimensional Lie algebra $\mathfrak{g}$. It is known that $\widehat{\mathfrak{g}}$ is a one-dimensional central extension of the *loop algebra* of the finite dimensional Lie algebra $\mathfrak{g}$. We denote this one-dimensional center by $\mathbb{R}\mathbf{c}$, where $\mathbf{c} \in \widehat{\mathfrak{h}}$. The Cartan subalgebra in the extended affine algebra $\widehat{\mathfrak{g}}^e$ can then be written as

$$\widehat{\mathfrak{h}}^e = \mathbb{R}\mathbf{c} \oplus \mathfrak{h} \oplus \mathbb{R}\mathbf{D}, \quad (2.24)$$

in which we have assumed (as we may) that $\mathfrak{h}$ coincides with the fixed Cartan subalgebra of $\mathfrak{g}$ chosen in section 2.1. The finite dimensional roots $\alpha \in \Delta$ can then be extended to elements of $(\widehat{\mathfrak{h}}^e)^*$ by stipulating that

$$\langle \alpha, \mathbf{c} \rangle = \langle \alpha, \mathbf{D} \rangle = 0 \quad \text{for each } \alpha \in \Delta. \quad (2.25)$$

In particular the element ρ defined in (2.4), which for us always represents an object from the classical group, extends to an element of $(\widehat{\mathfrak{h}}^e)^*$ that is trivial on $\mathbf{c}$ and $\mathbf{D}$. We may then identify the first ℓ simple roots of the affine Lie algebra with those of its classical counterpart:

$$a_i = \alpha_i \text{ for } i = 1, 2, \dots, \ell. \quad (2.26)$$

We likewise extend the fundamental weights ω_j for $j = 1, \dots, \ell$ to elements of $(\widehat{\mathfrak{h}}^e)^*$ by setting

$$\langle \omega_j, \mathbf{c} \rangle = \langle \omega_j, \mathbf{D} \rangle = 0. \quad (2.27)$$

Also, we can identify the affine coroots $h_1, h_2, \dots, h_\ell$ with the corresponding classical coroots defined in §2.1 under the same name. It remains to describe both $a_{\ell+1}$ and $h_{\ell+1}$ in terms of the underlying finite-dimensional root data.

Let $\iota \in \widehat{\Delta}$ be the minimal positive imaginary root. It is characterized as the unique linear map $\iota \in (\widehat{\mathfrak{h}}^e)^*$ satisfying the condition that

$$\begin{aligned} \langle \iota, X \rangle &= 0 \quad \text{for } X \in \mathbb{R}\mathbf{c} \oplus \mathfrak{h} \\ \text{and } \langle \iota, \mathbf{D} \rangle &= 1. \end{aligned} \quad (2.28)$$

We then have the following explicit description of the real and imaginary roots of $\widehat{\mathfrak{g}}^e$:

$$\widehat{\Delta}_W = \{ \alpha + n\iota \mid \alpha \in \Delta, n \in \mathbb{Z} \} \quad (2.29)$$

$$\widehat{\Delta}_I = \{ n\iota \mid n \neq 0, n \in \mathbb{Z} \}. \quad (2.30)$$

The classical roots Δ can be regarded as the subset of $\widehat{\Delta}_W$ with $n = 0$ in this parametrization. One then has

$$a_{\ell+1} = -\alpha_0 + \iota, \quad (2.31)$$

where we recall that α_0 denotes the highest root of the underlying finite-dimensional root system.

We shall now give a similar description of $h_{\ell+1}$. To do so, we recall from [22, (6.2.1)] that one can define a symmetric, non-degenerate, invariant bilinear form $(\cdot | \cdot)$ on $\widehat{\mathfrak{h}}^e$. The form is first defined on $\widehat{\mathfrak{h}}$ in terms of certain labels of the affine Dynkin diagram (coming from the coefficients of α_0 when expanded as a sum of the finite simple roots), and then is extended to all of $\widehat{\mathfrak{h}}^e$ by setting

$$\begin{aligned} (h_i | \mathbf{D}) &= 0, \quad \text{for } i = 1, \dots, \ell, \\ (h_{\ell+1} | \mathbf{D}) &= 1, \\ \text{and } (\mathbf{D} | \mathbf{D}) &= 0. \end{aligned} \quad (2.32)$$

The form $(\cdot | \cdot)$ also induces a symmetric bilinear form on $(\widehat{\mathfrak{h}}^e)^*$, which we continue to denote by $(\cdot | \cdot)$, and which has the following characterization:

$$\begin{aligned} (a_i | a_j) &= (\alpha_i, \alpha_j), \quad \text{for } i, j \in \{1, \dots, \ell\}, \\ (a_i | a_{\ell+1}) &= (\alpha_i, -\alpha_0), \quad \text{for } i \in \{1, \dots, \ell\}, \\ \text{and } (a_{\ell+1} | a_{\ell+1}) &= (\alpha_0, \alpha_0) = 2. \end{aligned} \quad (2.33)$$

It is easy to see that the above equations imply that

$$(\iota | \iota) = 0 \quad \text{and} \quad (\iota | a_i) = 0 \quad \text{for } i = 1, \dots, \ell. \quad (2.34)$$

Note that together with the relations

$$\begin{aligned} (\Lambda_{\ell+1} | \Lambda_{\ell+1}) &= 0, \\ (a_i | \Lambda_{\ell+1}) &= 0, \quad \text{for } i = 1, \dots, \ell, \\ \text{and } (a_{\ell+1} | \Lambda_{\ell+1}) &= 1, \end{aligned} \quad (2.35)$$

(2.33) and (2.34) completely specify the form $(\cdot|\cdot)$ on $(\widehat{\mathfrak{h}}^e)^*$ (and hence $\widehat{\mathfrak{h}}^e$). We define normalized coroots by the formula

$$h'_{a_i} := \frac{(a_i|a_i)}{2} h_i, \quad i \in \{1, \dots, \ell+1\}. \quad (2.36)$$

More generally for

$$b = \sum_{i=1}^{\ell+1} \kappa_i a_i \quad (2.37)$$

we set

$$h'_b := \sum_{i=1}^{\ell+1} \kappa_i h'_{a_i}. \quad (2.38)$$

Using (2.26) and (2.31) these conventions give a definition for h'_l .

For any real root b we have defined the corresponding coroot h_b in (2.23) in terms of the Weyl group action. One then has that

$$h'_b = \frac{(b|b)}{2} h_b \quad (2.39)$$

(see for example [22, (5.1.1)]). We now set

$$h_l := \frac{2}{(\alpha_0, \alpha_0)} h'_l = h'_l, \quad (2.40)$$

using $(\alpha_0, \alpha_0) = 2$. Suppose $a = \alpha + n\iota \in \widehat{\Delta}_W$ with $\alpha \in \Delta$ and $n \in \mathbb{Z}$ as in (2.29). Then in fact

$$h_a = \frac{(\alpha|\alpha)}{(a|a)} h_\alpha + \frac{2}{(a|a)} n h_l = h_\alpha + \frac{2}{(\alpha|\alpha)} n h_l, \quad (2.41)$$

where in the second equality we have used the fact that $(a|a) = (\alpha|\alpha)$, a consequence of (2.34). In particular we also have the formula

$$h_{\ell+1} = h_{-\alpha_0} + h_l, \quad (2.42)$$

in analogy to (2.31). One may check that h_l is also a generator for the one-dimensional center $\mathbb{R}\mathbf{c}$ of $\widehat{\mathfrak{g}}^e$ and that we have the direct sum decompositions

$$\widehat{\mathfrak{h}}^e = \mathbb{R}h_l \oplus \mathfrak{h} \oplus \mathbb{R}\mathbf{D} \quad \text{and} \quad \widehat{\mathfrak{h}} = \mathbb{R}h_l \oplus \mathfrak{h} \quad (2.43)$$

similar to (2.24). This decomposition of $\widehat{\mathfrak{h}}^e$ will be frequently used later in this paper.

C. Lemmas on the Affine Weyl Group

2.6. The affine Weyl group $\widehat{W}$ was defined above as the group generated by the reflections $w_1, \dots, w_{\ell+1}$ from (2.20). It also has a more classical description as the semidirect product

$$\widehat{W} = W \ltimes Q^\vee. \quad (2.44)$$

More concretely, the elements b of the coroot lattice $Q^\vee$ correspond to translations T_b in the Weyl group $\widehat{W}$. Recall that T_b fixes ι , as do all elements in $\widehat{W}$. The action of T_b on $\lambda \in \text{Span}(a_1, \dots, a_\ell)$ is given by the formula

$$T_b : \lambda \mapsto \lambda + \langle \lambda, b \rangle \iota. \quad (2.45)$$

It is also possible to obtain a general formula for the action of T_b on an element in $(\widehat{\mathfrak{h}}^e)^*$, though we will not need it here.

We shall now state a formula for the action of $\widehat{W}$ on $\widehat{\mathfrak{h}}^e$, which we shall make frequent use of later. The general element of $\widehat{\mathfrak{h}}^e$ can be written in terms of the decomposition (2.43) as $mh_{\mathfrak{t}} + h + r\mathbf{D}$. If we use (2.44) to factor $w \in \widehat{W}$ as $w = \widetilde{w}T_b$ for some $\widetilde{w} \in W$ and $b \in Q^\vee$, then we have

$$w \cdot (mh_{\mathfrak{t}} + h + r\mathbf{D}) = \left[-\frac{r(b,b)}{2} + (h,b) + m \right] h_{\mathfrak{t}} - r\widetilde{w}(b) + \widetilde{w}(h) + r\mathbf{D} \quad (2.46)$$

(see [8, p. 309]). Note that in particular $\widehat{W}$ fixes $h_{\mathfrak{t}}$, $\mathfrak{t}$, and $\mathbf{c}$. Combined with the fact that (2.23) is well-defined, this shows that the coroots satisfy

$$wh_a = h_{wa} \quad (2.47)$$

for any $w \in \widehat{W}$ and $a \in \widehat{\Delta}$.

2.7. For each $w \in \widehat{W}$ we denote by $\ell(w)$ the length of w , i.e., the minimal length of a word in $w_1, \dots, w_{\ell+1}$ which represents w . We then have the following estimate:

Lemma. *Let $\|\cdot\|$ be an arbitrary norm on $\mathfrak{h}$. Then there exist constants $E, E' > 0$ depending only on the (finite-dimensional) root system Δ and the choice of norm $\|\cdot\|$ such that*

$$E' \|b\| - \#(\Delta_+) \leq \ell(w) = \ell(w^{-1}) \leq E \|b\| + \#(\Delta_+), \quad (2.48)$$

for any element $w \in \widehat{W}$ of the form $w = T_b \widetilde{w}$ or $w = \widetilde{w} T_b$, where $b \in Q^\vee$ and $\widetilde{w} \in W$.

Proof. Assume that w has the form $T_b \widetilde{w}$ (the statement for $w = \widetilde{w} T_b$ is equivalent). From [21, Proposition 1.23] one has

$$\ell(w) = \sum_{\alpha \in \Delta_+} |\langle \alpha, b \rangle - \chi_{\Delta_-}(\widetilde{w}^{-1} \alpha)|, \quad (2.49)$$

where χ_{Δ_-} is the characteristic function of Δ_- . By the triangle inequality

$$\ell(w) \leq \sum_{\alpha \in \Delta_+} (|\langle \alpha, b \rangle| + 1) = \sum_{\alpha \in \Delta_+} |\langle \alpha, b \rangle| + \#(\Delta_+). \quad (2.50)$$

Similarly, we deduce from (2.49) that

$$\ell(w) \geq \sum_{\alpha \in \Delta_+} (|\langle \alpha, b \rangle| - 1) = \sum_{\alpha \in \Delta_+} |\langle \alpha, b \rangle| - \#(\Delta_+). \quad (2.51)$$

Writing $b = \sum_{i=1}^{\ell} d_i \omega_i^\vee$ in terms of the fundamental coweights $\omega_i^\vee$ and letting

$$M = \max_{\substack{\alpha \in \Delta_+ \\ 1 \leq i \leq \ell}} \langle \alpha, \omega_i^\vee \rangle, \quad (2.52)$$

we have $|\langle \alpha, b \rangle| \leq M \sum_{i=1}^{\ell} |d_i|$ for any $\alpha \in \Delta_+$. On the other hand,

$$\sum_{\alpha \in \Delta_+} |\langle \alpha, b \rangle| \geq \sum_{\alpha \in \Pi} |\langle \alpha, b \rangle| = \sum_{i=1}^{\ell} |\langle \alpha_i, b \rangle| = \sum_{i=1}^{\ell} |d_i|. \quad (2.53)$$

As all norms on a finite dimensional vector space are equivalent, the assertions of the lemma follow. $\square$

2.8. For any element $w \in \widehat{W}$ we set

$$\widehat{\Delta}_w = \{a \in \widehat{\Delta}_+ \mid w^{-1}a < 0\}. \quad (2.54)$$

It is a standard result that

$$\#(\widehat{\Delta}_w) = \ell(w) = \ell(w^{-1}). \quad (2.55)$$

Moreover, given a reduced decomposition $w = w_{i_r} \dots w_{i_1}$, $r = \ell(w)$, the set $\widehat{\Delta}_w$ can be described as

$$\widehat{\Delta}_w = \{\beta_1, \dots, \beta_r\}, \quad (2.56)$$

where $\beta_j = w_{i_r} \dots w_{i_{j+1}}(a_{i_j})$ for $j = 1, \dots, r$. We also set

$$\widehat{\Delta}_{-,w} = \{a \in \widehat{\Delta}_- \mid wa > 0\} = w^{-1}\widehat{\Delta}_w. \quad (2.57)$$

Similarly to (2.56) it equals

$$\widehat{\Delta}_{-,w} = \{\gamma_1, \dots, \gamma_r\}, \quad (2.58)$$

where $\gamma_j = w^{-1}\beta_j = w_{i_1} \dots w_{i_j}(a_{i_j}) = -w_{i_1} \dots w_{i_{j-1}}(a_{i_j})$ for $j = 1, \dots, r$.

2.9. Considering the classical Weyl group W as the subgroup of $\widehat{W}$ generated by $w_1, \dots, w_\ell$, let W^θ denote the *Kostant coset representatives* for the quotient $W \backslash \widehat{W}$:

$$W^\theta = \{w \in \widehat{W} \mid w^{-1}\alpha_i > 0 \text{ for } i = 1, \dots, \ell\}. \quad (2.59)$$

For any $i = 1, \dots, \ell$ and $w \in W^\theta$ let us write

$$w^{-1}\alpha_i = \sigma_i + \kappa_i(w^{-1})\iota, \quad \text{where } \sigma_i \in \Delta \text{ and } \kappa_i(w^{-1}) \in \mathbb{Z}_{\geq 0}, \quad (2.60)$$

which is possible since the Weyl translates of the α_i lie in $\widehat{\Delta}_w$ (see (2.29)).

Lemma. For any $w \in W^\theta$ and $i = 1, \dots, \ell$, we have that

$$\kappa_i(w^{-1}) \leq \ell(w) + 1 = \ell(w^{-1}) + 1. \quad (2.61)$$

Proof. Suppose first that σ_i in (2.60) is positive. Then

$$w^{-1}(-\alpha_i + n\iota) = -\sigma_i + (n - \kappa_i(w^{-1}))\iota \quad (2.62)$$

since $\widehat{W}$ preserves the imaginary root ι . The root $-\alpha_i + n\iota$ is positive if $n > 0$, while $w^{-1}(-\alpha_i + n\iota)$ is negative if $n - \kappa_i(w^{-1}) \leq 0$. Hence $\widehat{\Delta}_w$ from (2.54) includes the roots

$$-\alpha_i + \iota, -\alpha_i + 2\iota, \dots, -\alpha_i + \kappa_i(w^{-1})\iota. \quad (2.63)$$

From this and (2.55) we conclude that $\kappa_i(w^{-1}) \leq \ell(w^{-1}) = \ell(w)$. A similar analysis for the case of $\sigma_i < 0$ shows $\widehat{\Delta}_w$ contains the string

$$-\alpha_i + \iota, \dots, -\alpha_i + (\kappa_i(w^{-1}) - 1)\iota, \quad (2.64)$$

and so $\kappa_i(w^{-1}) \leq \ell(w^{-1}) + 1$. □

D. Loop Groups and Some Decompositions

2.10. We next introduce some notation related to loop groups following [6], which we briefly review here. Let $\mathfrak{g}_{\mathbb{Z}} \subset \mathfrak{g}$ and $\widehat{\mathfrak{g}}_{\mathbb{Z}}^e \subset \widehat{\mathfrak{g}}^e$ be Chevalley $\mathbb{Z}$ -forms of the Lie algebras defined above. Given a dominant integral weight $\lambda \in \widehat{\mathfrak{h}}^*$ extended to $(\widehat{\mathfrak{h}}^e)^*$ as in (2.16-2.17), let V^λ be the corresponding irreducible highest-weight representation of $\widehat{\mathfrak{g}}^e$. Also, let $V_{\mathbb{Z}}^\lambda \subset V^\lambda$ denote a Chevalley $\mathbb{Z}$ -form for this representation, chosen compatibly with the Chevalley forms $\mathfrak{g}_{\mathbb{Z}}$ and $\widehat{\mathfrak{g}}_{\mathbb{Z}}^e$ and with suitable divided powers as in [5].

For any commutative ring R with unit, let $\mathfrak{g}_R$, $\widehat{\mathfrak{g}}_R^e$, and V_R^λ be the objects obtained by tensoring the respective objects $\mathfrak{g}_{\mathbb{Z}}$, $\widehat{\mathfrak{g}}_{\mathbb{Z}}^e$, and $V_{\mathbb{Z}}^\lambda$ over $\mathbb{Z}$ with R . For any field k , we let $\widehat{G}_k^\lambda \subset \text{Aut}(V_k^\lambda)$ be the group defined in [6, (7.21)] by exponentiation of the corresponding Lie algebra representation. When no subscript is present, we shall implicitly assume that $k = \mathbb{R}$, so that for example $\widehat{G}^\lambda$ is taken to mean $\widehat{G}_{\mathbb{R}}^\lambda$, etc.

For $a \in \widehat{\Delta}_W$ and $u \in k$ we define $\chi_a(u)$ as in [6, (7.14)], which parameterizes the one-parameter root group corresponding to a . Using $\chi_a(u)$ we may then define the elements

$$w_a(s) := \chi_a(s) \chi_{-a}(-s^{-1}) \chi_a(s) \quad (2.65)$$

and

$$h_a(s) := w_a(s) w_a(1)^{-1} \quad (2.66)$$

for $s \in k^*$. We shall use the abbreviation $h_i(s) = h_{a_i}(s)$ for $i = 1, \dots, \ell + 1$. We shall fix λ throughout and shorten our notation to $\widehat{G} := \widehat{G}_{\mathbb{R}}^\lambda$ and make a similar convention for $\widehat{G}_k$.

In case $k = \mathbb{R}$, we let $\{\cdot, \cdot\}$ denote the real, positive-definite inner product on $V_{\mathbb{R}}^\lambda$ (see [6, §16] and the references therein). We then define subgroups

$$\begin{aligned} \widehat{\Gamma} &= \left\{ \gamma \in \widehat{G} \mid \gamma V_{\mathbb{Z}}^\lambda = V_{\mathbb{Z}}^\lambda \right\} \\ \text{and } \widehat{K} &= \left\{ k \in \widehat{G} \mid \{k\xi, k\eta\} = \{\xi, \eta\} \text{ for } \xi, \eta \in V_{\mathbb{R}}^\lambda \right\}. \end{aligned} \quad (2.67)$$

Fix a coherently ordered basis $\mathcal{B}$ of $V_{\mathbb{Z}}^\lambda$ in the sense of [6, p. 60], and let $\widehat{H}$, $\widehat{U}$, and $\widehat{B}$ respectively denote the subgroups of $\widehat{G}$ consisting of diagonal, unipotent upper triangular, and upper triangular matrices with respect to $\mathcal{B}$. The subgroup $\widehat{U}$ contains the one-parameter subgroups $\chi_a(u)$ for each $a \in \widehat{\Delta}_+$. The subgroup $\widehat{H}$ normalizes $\widehat{U}$ and

$$\widehat{B} = \widehat{H} \ltimes \widehat{U} \quad (2.68)$$

is their semi-direct product. The group $\widehat{H}$ is isomorphic to $(\mathbb{R}^*)^{\ell+1}$ via the map

$$(s_1, \dots, s_{\ell+1}) \mapsto h_1(s_1) \cdots h_{\ell+1}(s_{\ell+1}). \quad (2.69)$$

We use this parametrization to define the map $h \mapsto h^{a_i}$ on $\widehat{H}$ for any simple root a_i , via the formula $h_j(s)^{a_i} = s^{\langle a_i, h_j \rangle}$. The products on the righthand side of (2.69) with each $s_i > 0$ form a subgroup $\widehat{A} \cong \mathbb{R}_{>0}^{\ell+1}$ of $\widehat{H}$, so that $\widehat{A}$ is isomorphic to $\widehat{\mathfrak{h}}$ by the logarithm map

$$\ln : h_1(s_1) \cdots h_{\ell+1}(s_{\ell+1}) \mapsto \ln(s_1) h_1 + \cdots + \ln(s_{\ell+1}) h_{\ell+1}. \quad (2.70)$$

Both $\widehat{H}$ and $\widehat{A}$ have Lie algebra $\widehat{\mathfrak{h}}$, and in fact

$$\widehat{H} = \widehat{A} \times (\widehat{H} \cap \widehat{K}) \quad (2.71)$$

holds as a direct product decomposition. The notation (2.7) extends to give a character $a \mapsto a^\lambda$ of $\widehat{A}$ for any $\lambda \in (\widehat{\mathfrak{h}}^e)^*$. For any linear combination

$$X = c_1 a_1 + \cdots + c_{\ell+1} a_{\ell+1}, \quad c_i \in \mathbb{Z}, \quad (2.72)$$

we define the map

$$h_X(s) := h_1(s)^{c_1} \cdots h_{\ell+1}(s)^{c_{\ell+1}} \quad \text{for } s \in \mathbb{R}^*. \quad (2.73)$$

Using the relation $\iota = \alpha_0 + a_{\ell+1}$ from (2.31), this then allows us to define $h_\iota(s)$ for $s \in \mathbb{R}^*$.

Define

$$\begin{aligned} \widehat{H}_{cen} &:= \{h_\iota(s) \mid s \in \mathbb{R}^*\} \\ \text{and } \widehat{H}_{cl} &:= \left\{ \prod_{i=1}^{\ell} h_i(s_i) \mid s_1, \dots, s_\ell \in \mathbb{R}^* \right\}, \end{aligned} \quad (2.74)$$

and then set $\widehat{A}_{cen} = \widehat{A} \cap \widehat{H}_{cen}$ and $\widehat{A}_{cl} = \widehat{A} \cap \widehat{H}_{cl}$. We then have a direct product decomposition

$$\widehat{A} = \widehat{A}_{cen} \times \widehat{A}_{cl}. \quad (2.75)$$

Indeed, it suffices to show that $\widehat{A}_{cl} \cap \widehat{A}_{cen} = \{1\}$. Indeed, any element $x \in \widehat{A}_{cl} \cap \widehat{A}_{cen}$ can be written as

$$x = h_\iota(s) = h, \quad \text{with } s \in \mathbb{R}_{>0} \text{ and } h \in \widehat{A}_{cl}. \quad (2.76)$$

Recall from above that the group $\widehat{G}$ is defined with respect to a dominant integral weight λ . Because of [6, Proposition 20.2], there exists a positive integer m together with a homomorphism $\pi(\lambda, m\Lambda_{\ell+1}) : \widehat{G}^\lambda \hookrightarrow \widehat{G}^{m\Lambda_{\ell+1}}$. According to [6, p. 107] this map identifies the elements $h_i(s)$ in the two groups, for any $1 \leq i \leq \ell+1$. From (2.18) and (2.42) we see that the fundamental weight $\Lambda_{\ell+1}$ satisfies $\langle \Lambda_{\ell+1}, h_i \rangle = 1$ and $\langle \Lambda_{\ell+1}, h_i \rangle = 0$ for $i = 1, \dots, \ell$. In the group $\widehat{G}^{m\Lambda_{\ell+1}}$, which acts on the highest weight representation $V^{m\Lambda_{\ell+1}}$ with highest weight vector $v_{m\Lambda_{\ell+1}}$, [6, Lemma 11.2] shows

$$h_\iota(s) v_{m\Lambda_{\ell+1}} = s^{m\langle \Lambda_{\ell+1}, h_\iota \rangle} v_{m\Lambda_{\ell+1}} = s^m v_{m\Lambda_{\ell+1}}. \quad (2.77)$$

However, using (2.76) the same lemma shows $h_\iota(s) v_{m\Lambda_{\ell+1}} = h v_{m\Lambda_{\ell+1}} = v_{m\Lambda_{\ell+1}}$ because $h \in \widehat{A}_{cl}$. Thus $s^m = 1$ from which we can conclude that $s = 1$ itself.

2.11. For any field k , we define $\widehat{H}_k$ to be the subgroup of $\widehat{G}_k$ consisting of diagonal matrices with respect to the coherent basis $\mathcal{B}$. Similarly we set $\widehat{B}_k$ and $\widehat{U}_k$ to be the groups of upper triangular and unipotent upper triangular matrices with respect to $\mathcal{B}$, respectively. As in (2.68), $\widehat{B}_k$ is the semi-direct product of $\widehat{H}_k$ and $\widehat{U}_k$. Moreover, the group $\widehat{H}_k$ is isomorphic to $(k^*)^{\ell+1}$ via (2.69) (recall that the elements $h_a(s)$ from (2.66) are defined for $s \in k^*$).

The group $\widehat{G}_k$ is equipped with a Tits system (see [6, §13-14]) which identifies the affine Weyl group $\widehat{W}$ from section 2.6 as the quotient $N(\widehat{H}_k)/\widehat{H}_k = \widehat{W}$, where $N(\widehat{H}_k)$ is the normalizer of $\widehat{H}_k$ in $\widehat{G}$. Thus each $w \in \widehat{W}$ has a representative in $N(\widehat{H}_k)$, which we continue to denote by w . Moreover,

if w is written as a word in the generators $w_1, \dots, w_{\ell+1}$ from (2.20), a representative is furnished by the corresponding word in the elements $w_{a_1}(1), \dots, w_{a_{\ell+1}}(1)$ from (2.65). We shall tacitly identify each $w \in \widehat{W}$ with this particular representative. In the special case $k = \mathbb{R}$ these representatives lie in $\widehat{K}$. With these conventions, there exists a Bruhat decomposition

$$\widehat{G}_k = \bigcup_{w \in \widehat{W}} \widehat{B}_k w \widehat{B}_k, \quad (2.78)$$

where in fact each double coset $\widehat{B}_k w \widehat{B}_k$ is independent of the chosen representative w . Because of (2.68) and the fact that $\widehat{W}$ normalizes $\widehat{H}_k$, every element $g \in \widehat{G}_k$ can be written as

$$g = u_1 z w u_2, \quad \text{where } u_1, u_2 \in \widehat{U}_k, \quad w \in \widehat{W}, \quad \text{and } z \in \widehat{H}_k. \quad (2.79)$$

If $k = \mathbb{R}$, the elements $w \in \widehat{W}$ and $z \in \widehat{H}$ are uniquely determined by g , though u_1 and $u_2 \in \widehat{U}$ are not in general.

2.12. Next, we recall from [6, §16] that there exist Iwasawa decompositions

$$\widehat{G} = \widehat{U} \widehat{A} \widehat{K} = \widehat{K} \widehat{A} \widehat{U} \quad (2.80)$$

with respect to the subgroups $\widehat{U}$, $\widehat{A}$, and $\widehat{K}$ defined in section 2.10, with uniqueness of decomposition in either particular fixed order. Given an element $g \in \widehat{G}$ we denote by $\text{Iw}_{\widehat{A}}(g) \in \widehat{A}$ its projection onto the $\widehat{A}$ factor in the first of the above decompositions. Note that $\text{Iw}_{\widehat{A}} : \widehat{G} \rightarrow \widehat{A}$ is left $\widehat{U}$ -invariant and right $\widehat{K}$ -invariant by construction.

For $r \in \mathbb{R}$, write $s = e^r$ and define the exponentiated degree operator on $V_{\mathbb{R}}^{\lambda}$ by the formula

$$\eta(s) = \exp(r\mathbf{D}). \quad (2.81)$$

The element $\eta(s)$ acts on the one-parameter subgroups $\chi_{\alpha+nl}(\cdot)$ by

$$\eta(s) \chi_{\alpha+nl}(u) \eta(s)^{-1} = \chi_{\alpha+nl}(s^n u). \quad (2.82)$$

It furthermore acts as a diagonal operator with respect to the coherently ordered basis $\mathcal{B}$, and consequently it normalizes $\widehat{U}$ and commutes with $\widehat{A}$. It then follows that

$$\eta(s) \widehat{G} = \{\eta(s)g \mid g \in \widehat{G}\} = \widehat{U} \eta(s) \widehat{A} \widehat{K}. \quad (2.83)$$

We then extend the function $\text{Iw}_{\widehat{A}}$ above to a function $\text{Iw}_{\eta(s)\widehat{A}} : \eta(s)\widehat{G} \rightarrow \eta(s)\widehat{A}$ by defining

$$\text{Iw}_{\eta(s)\widehat{A}}(\eta(s)g) = \eta(s) \text{Iw}_{\widehat{A}}(g) \in \eta(s)\widehat{A}. \quad (2.84)$$

This extension is obviously right $\widehat{K}$ -invariant, and is also left $\widehat{U}$ -invariant because $\eta(s)$ normalizes $\widehat{U}$. It furthermore satisfies

$$\text{Iw}_{\eta(s)\widehat{A}}(\eta(s)a_1 u a_2 k) = \text{Iw}_{\eta(s)\widehat{A}}(\eta(s)a_1) a_2 = \eta(s)a_1 a_2 \quad (2.85)$$

for any $u \in \widehat{U}$, $a_1, a_2 \in \widehat{A}$, and $k \in \widehat{K}$.

We extend the logarithm map $\ln : \widehat{A} \cong \widehat{\mathfrak{h}}$ defined in (2.70) to $\eta(s)\widehat{A}$ by the rule

$$\ln(\eta(s)a) = r\mathbf{D} + \ln(a) \quad \text{for } a \in \widehat{A} \quad (2.86)$$

(cf. (2.81)). For the duration of the paper we make the important restriction to consider *only* the case that $r > 0$.¹

¹Note that the convention in [6] is to also consider $r > 0$, although $\eta(s)$ from (2.81) is instead parameterized there as $\eta(s) = \exp(-r\mathbf{D})$. This is because here we work with the Iwasawa decomposition (2.80) having $\widehat{K}$ on the right, whereas in *op. cit.* $\widehat{K}$ is on the left.

E. Adelic Loop Groups

2.13. In this section, we review some aspects of adelic loop groups and their decompositions; further details can be found in [8, 11]. Let $\mathcal{V}$ denote the set of finite places of $\mathbb{Q}$, each of which can be identified with a prime number p and the p -adic norm $|\cdot|_p$. For each $p \in \mathcal{V}$ the field $\mathbb{Q}_p$ is the corresponding completion of $\mathbb{Q}$ and has ring of integers $\mathbb{Z}_p$. We write $\mathcal{V}^e = \mathcal{V} \cup \{\infty\}$ for the set of all places of $\mathbb{Q}$, where the place ∞ corresponds to the archimedean valuation $|\cdot|_\infty$, i.e., the usual absolute value on $\mathbb{Q}_\infty = \mathbb{R}$. The adeles are defined as the ring $\mathbb{A} = \prod'_{p \in \mathcal{V}^e} \mathbb{Q}_p$, where the prime indicates the restricted direct product of the factors with respect to the $\mathbb{Z}_p$. Likewise, the finite adeles $\mathbb{A}_f$ are the restricted direct product of all $\mathbb{Q}_p$, $p \in \mathcal{V}$, with respect to the $\mathbb{Z}_p$. For each $a = (a_p) \in \mathbb{A}$ the adelic valuation is defined as $|a|_\mathbb{A} := \prod_{p \in \mathcal{V}^e} |a_p|_p$. We also write $\mathbb{I} = \mathbb{A}^*$ for the group of ideles and $\mathbb{I}_f = \mathbb{I} \cap \mathbb{A}_f$ the group of finite ideles.

In section 2.10 we introduced the exponentiated group $\widehat{G}_k$ for any field k , in particular $k = \mathbb{Q}_p$ for any $p \in \mathcal{V}^e$. For shorthand denote $\widehat{G}_p := \widehat{G}_{\mathbb{Q}_p}$, so that $\widehat{G}_\infty$ is just the real group $\widehat{G}$. For each $p \in \mathcal{V}$, we set

$$\widehat{K}_p = \{g \in \widehat{G}_p \mid gV_{\mathbb{Z}_p}^\lambda = V_{\mathbb{Z}_p}^\lambda\}. \quad (2.87)$$

By convention, we also set $\widehat{K}_\infty$ to be the group $\widehat{K}$ introduced in (2.67). The adelic loop group is then defined as

$$\widehat{G}_\mathbb{A} := \prod'_{p \in \mathcal{V}^e} \widehat{G}_p, \quad (2.88)$$

where the product is restricted with respect to the family of subgroups $\{\widehat{K}_p\}_{p \in \mathcal{V}}$. We also define

$$\widehat{K}_\mathbb{A} = \prod_{p \in \mathcal{V}^e} \widehat{K}_p \subset \widehat{G}_\mathbb{A}. \quad (2.89)$$

Analogously, the groups $\widehat{G}_{\mathbb{A}_f}$ and $\widehat{K}_{\mathbb{A}_f}$ are defined by replacing $\mathcal{V}^e$ with $\mathcal{V}$ in (2.88) and (2.89), respectively.

We set $\widehat{H}_p := \widehat{H}_{\mathbb{Q}_p} \subset \widehat{G}_p$ to be the group defined at the beginning of §2.11, where we remarked that it is generated by the elements $h_1(s), h_2(s), \dots, h_{\ell+1}(s)$ for $s \in \mathbb{Q}_p^*$; thus for example, $\widehat{H}_\infty = \widehat{H}$. Define $\widehat{H}_\mathbb{A} = \prod'_{p \in \mathcal{V}^e} \widehat{H}_p$, where the product is restricted with respect to the family of subgroups $\{\widehat{H}_p \cap \widehat{K}_p\}_{p \in \mathcal{V}}$. Analogously to (2.69), every element $h \in \widehat{H}_\mathbb{A}$ has an expression

$$h = \prod_{i=1}^{\ell+1} h_i(s_i), \quad \text{where each } s_i \in \mathbb{I}. \quad (2.90)$$

For such an expression, we define its norm to be the element of $\widehat{A}$ given by the product

$$|h| = \prod_{i=1}^{\ell+1} h_i(|s_i|_\mathbb{A}), \quad (2.91)$$

which can be shown to be uniquely determined by $h \in \widehat{H}_\mathbb{A}$ independently of its factorization (2.90). We set $\widehat{U}_\mathbb{A} = \prod'_{p \in \mathcal{V}^e} \widehat{U}_{\mathbb{Q}_p}$, where the product is restricted with respect to the family $\{\widehat{U}_{\mathbb{Q}_p} \cap \widehat{K}_p\}_{p \in \mathcal{V}}$.

We shall also write $\widehat{B}_{\mathbb{A}} = \widehat{U}_{\mathbb{A}} \cdot \widehat{H}_{\mathbb{A}}$, which itself is the restricted direct product of all $\widehat{U}_{\mathbb{Q}_p} \cdot \widehat{H}_p$ with respect to their intersections with $\widehat{K}_p$.

2.14. We next state the adelic analogues of the Iwasawa decompositions (2.80) and (2.83). First, for each $p \in \mathcal{V}$ there is the p -adic Iwasawa decomposition $\widehat{G}_{\mathbb{Q}_p} = \widehat{U}_{\mathbb{Q}_p} \widehat{H}_{\mathbb{Q}_p} \widehat{K}_{\mathbb{Q}_p}$, which is not a direct product decomposition because $\widehat{H}_{\mathbb{Q}_p} \cap \widehat{K}_{\mathbb{Q}_p}$ is nontrivial. Together with the $p = \infty$ decomposition (2.80), these local decompositions give the adelic Iwasawa decomposition

$$\widehat{G}_{\mathbb{A}} = \widehat{U}_{\mathbb{A}} \widehat{H}_{\mathbb{A}} \widehat{K}_{\mathbb{A}}. \quad (2.92)$$

Although the adelic Iwasawa factorization

$$g = u_g h_g k_g, \quad u_g \in \widehat{U}_{\mathbb{A}}, h_g \in \widehat{H}_{\mathbb{A}}, k_g \in \widehat{K}_{\mathbb{A}}, \quad (2.93)$$

is not in general unique, the element $|h_g|$ defined using (2.91) is independent of it. We shall write the projection onto this element as the map

$$\begin{aligned} \text{Iw}_{\widehat{A}}^{\mathbb{A}} : \widehat{G}_{\mathbb{A}} &\rightarrow \widehat{A}, \\ g &\mapsto |h_g| \end{aligned} \quad (2.94)$$

onto the Iwasawa $\widehat{A}$ -factor of $\widehat{G}$. Note that $|h_g|$ is an element of the real group $\widehat{G}$; we do not adelize $\widehat{A}$.

Recall that we have defined $\eta(s)$ in (2.81) in the context of real groups, where it has a nontrivial action on $\widehat{G} = \widehat{G}_{\mathbb{Q}_{\infty}}$ by conjugation. After extending this action trivially to each $\widehat{G}_{\mathbb{Q}_p}$, $p \in \mathcal{V}$, there is then a twisted Iwasawa decomposition

$$\eta(s) \widehat{G}_{\mathbb{A}} = \widehat{U}_{\mathbb{A}} \eta(s) \widehat{H}_{\mathbb{A}} \widehat{K}_{\mathbb{A}}. \quad (2.95)$$

Moreover, if we write $\eta(s)g \in \eta(s)\widehat{G}_{\mathbb{A}}$ with respect to the above decomposition as

$$\eta(s)g = u_g \eta(s) h_g k_g, \quad u_g \in \widehat{U}_{\mathbb{A}}, h_g \in \widehat{H}_{\mathbb{A}}, k_g \in \widehat{K}_{\mathbb{A}}, \quad (2.96)$$

then the element $|h_g|$ from (2.91) is again uniquely determined. We shall denote this projection by

$$\begin{aligned} \text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}} : \eta(s)\widehat{G}_{\mathbb{A}} &\rightarrow \eta(s)\widehat{A}, \\ g &\mapsto \eta(s)|h_g|, \end{aligned} \quad (2.97)$$

generalizing (2.84).

2.15. In §2.10 the group $\widehat{G}_{\mathbb{Q}}$ was defined over the field $k = \mathbb{Q}$. It has embeddings $\widehat{G}_{\mathbb{Q}} \hookrightarrow \widehat{G}_{\mathbb{Q}_p}$ for each $p \in \mathcal{V}^e$, and hence a diagonal embedding

$$i : \widehat{G}_{\mathbb{Q}} \hookrightarrow \prod_{p \in \mathcal{V}^e} \widehat{G}_{\mathbb{Q}_p}. \quad (2.98)$$

Note that the righthand side is the direct product, not the restricted direct product: this is because $i(\widehat{G}_{\mathbb{Q}})$ is actually not contained in $\widehat{G}_{\mathbb{A}}$ (see [8, §2])². Define

$$\widehat{\Gamma}_{\mathbb{Q}} := i^{-1}(\widehat{G}_{\mathbb{A}}) \subset \widehat{G}_{\mathbb{Q}} \quad (2.99)$$

as the subgroup of $\widehat{G}_{\mathbb{Q}}$ which does embed into $\widehat{G}_{\mathbb{A}}$. We shall generally follow the common convention of identifying $\widehat{\Gamma}_{\mathbb{Q}}$ with its diagonally embedded image $i(\widehat{\Gamma}_{\mathbb{Q}})$.

²In contrast to the finite-dimensional situation, an element of $\widehat{G}_{\mathbb{Q}}$ can involve different prime denominators in each of infinitely many root spaces.

3 Iwasawa Inequalities

3.1. In (2.54) we associated to each $w \in \widehat{W}$ the finite set of roots

$$\widehat{\Delta}_{w^{-1}} = \{a \in \widehat{\Delta} \mid a > 0, wa < 0\}. \quad (3.1)$$

Recall that Weyl group fixes all imaginary roots, so $\widehat{\Delta}_{w^{-1}} \subset \widehat{\Delta}_W$. For each root $a \in \widehat{\Delta}_W$ let U_a denote the corresponding root group consisting of the elements $\{\chi_a(s) \mid s \in \mathbb{R}\}$, and fix any order on $\widehat{\Delta}$. Let $\widehat{U}_-$ be the subgroup of $\widehat{G}$ which acts by unipotent lower triangular matrices on the coherent basis $\mathcal{B}$ from section 2.10. We then have the *subgroups*

$$U_w = \prod_{a \in \widehat{\Delta}_{w^{-1}}} U_a = \widehat{U} \cap w^{-1} \widehat{U}_- w \subset \widehat{U} \quad (3.2)$$

and

$$U_{-,w} = w U_w w^{-1} = \prod_{\gamma \in \widehat{\Delta}_{-,w^{-1}}} U_\gamma = \widehat{U}_- \cap w \widehat{U} w^{-1} \subset \widehat{U}_-, \quad (3.3)$$

where the product is taken with respect to the fixed order and one has uniqueness of expression (cf. [7, Lemma 6.4 and Corollary 6.5]).

Similarly, each U_a has a rational subgroup $U_{a,\mathbb{Q}} = \{\chi_a(s) \mid s \in \mathbb{Q}\}$. Rational subgroups $U_{w,\mathbb{Q}} \subset U_w$ and $U_{-,w,\mathbb{Q}} \subset U_{-,w}$ are defined as products of $U_{a,\mathbb{Q}}$ and $U_{\gamma,\mathbb{Q}}$ over the roots a and γ appearing in (3.2) and (3.3), respectively. Because $\chi_a(s)$ is also defined for $s \in \mathbb{Q}_p$, we likewise have subgroups $U_{a,\mathbb{Q}_p}$, $U_{w,\mathbb{Q}_p}$, and $U_{-,w,\mathbb{Q}_p}$, and their adelic variants $U_{w,\mathbb{A}}$ and $U_{-,w,\mathbb{A}}$ defined as restricted direct products.

3.2. Recall the notation for the Iwasawa decomposition and its adelic variant introduced in (2.84), (2.94), and (2.97). For elements x, y in any group we shall use the shorthand notation $x^y := yxy^{-1}$.

Lemma. *i) Let $\gamma_{\mathbb{Q}} \in \widehat{\Gamma}_{\mathbb{Q}}$, regarded as diagonally embedded in $\widehat{G}_{\mathbb{A}}$ as in §2.15, and let $g \in \widehat{G}$. Use (2.79) to write $\gamma_{\mathbb{Q}} = u_1 z w u_2$ with $u_1, u_2 \in \widehat{U}_{\mathbb{Q}}$, $w \in \widehat{W}$, and $z \in \widehat{H}_{\mathbb{Q}}$. Regard the element $\gamma_{\mathbb{Q}} \eta(s) g \in \eta(s) \widehat{G}_{\mathbb{A}}$ and define the element $\text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}(\gamma_{\mathbb{Q}} \eta(s) g) \in \eta(s) \widehat{A}$ as in (2.97). Then*

$$\text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}(\gamma_{\mathbb{Q}} \eta(s) g) = \text{Iw}_{\eta(s)\widehat{A}}(\eta(s) g)^w \cdot \text{Iw}_{\widehat{A}}^{\mathbb{A}}(w u_w) \quad (3.4)$$

for some element $u_w \in U_{w,\mathbb{A}}$ (depending on $\gamma_{\mathbb{Q}}$ and $\eta(s) g$).

ii) If $\gamma \in \widehat{\Gamma} \subset \widehat{G}$,

$$\text{Iw}_{\eta(s)\widehat{A}}(\gamma \eta(s) g) = \text{Iw}_{\eta(s)\widehat{A}}(\eta(s) g)^w \cdot \text{Iw}_{\widehat{A}}^{\mathbb{A}}(w u_w) \quad (3.5)$$

for some element $u_w \in U_{w,\mathbb{A}}$ (again depending on γ and $\eta(s) g$).

Proof. First we note that since $z \in \widehat{H}_{\mathbb{Q}}$ we have that $\text{Iw}_{\widehat{A}}^{\mathbb{A}}(z) = 1$. Using the left $\widehat{U}_{\mathbb{A}}$ -invariance, the commutativity of $\widehat{A}$ with $\eta(s)$, and the right $\widehat{K}_{\mathbb{A}}$ -invariance of $\text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}$, we compute

$$\begin{aligned} \text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}(\gamma_{\mathbb{Q}} \eta(s) g) &= \text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}(u_1 z w u_2 \eta(s) g) \\ &= \text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}(z w u_2 \eta(s) g) \\ &= \text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}(w u_2 u' \eta(s) h'), \end{aligned} \quad (3.6)$$

where

$$\eta(s)g = u' \eta(s) h' k', \quad \text{with } u' \in \widehat{U}, h' = \text{Iw}_{\widehat{A}}(g) \in \widehat{A}, \text{ and } k' \in \widehat{K}, \quad (3.7)$$

is the (real, and hence also adelic) Iwasawa decomposition of $\eta(s)g$. Let $u = u_2 u' \in \widehat{U}_{\mathbb{A}}$. Again using these properties plus (2.85) we then have

$$\begin{aligned} \text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}(\gamma_{\mathbb{Q}} \eta(s)g) &= \text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}(wu \eta(s)h') \\ &= \text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}(w \eta(s)h' w^{-1} \cdot w \cdot (\eta(s)h')^{-1} \cdot u \cdot (\eta(s)h')) \\ &= \text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}((\eta(s)h')^w \cdot w u'') \\ &= (\eta(s)h')^w \text{Iw}_{\widehat{A}}^{\mathbb{A}}(w u''), \end{aligned} \quad (3.8)$$

where $u'' = (\eta(s)h')^{-1}u(\eta(s)h')$ lies in $\widehat{U}_{\mathbb{A}}$ because $\eta(s)$ and $h' \in \widehat{A}$ both normalize $\widehat{U}_{\mathbb{A}}$. Furthermore, u'' can be factored as $u'' = u''' u_w$, where $w u''' w^{-1} \in \widehat{U}_{\mathbb{A}}$ and $u_w \in U_{w, \mathbb{A}}$. Thus $w u'' = (w u''' w^{-1}) w u_w$ and formula (3.4) now follows from once more applying the left $\widehat{U}_{\mathbb{A}}$ -invariance of $\text{Iw}_{\eta(s)\widehat{A}}^{\mathbb{A}}$. This proves part i).

To prove part ii), let $\gamma_{\mathbb{Q}} \in \widehat{\Gamma}_{\mathbb{Q}}$ be the diagonal embedding of γ into $\widehat{G}_{\mathbb{A}}$. Since $\eta(s)g \in \widehat{G}$, the archimedean component of $\gamma_{\mathbb{Q}} \eta(s)g$ is $\gamma \eta(s)g$. At the same time, its nonarchimedean components are γ_p . These lie in $\widehat{K}_p$ since they preserve the lattice $V_{\mathbb{Z}_p}^{\lambda}$ (as a consequence of the defining fact that $\gamma \in \widehat{G}$ preserves $V_{\mathbb{Z}}^{\lambda}$). Hence the projection of $\gamma_{\mathbb{Q}} \eta(s)g$ onto $\widehat{G}_{\mathbb{A}_f}$ lies in $\widehat{K}_{\mathbb{A}_f}$, and by definition (2.94) the left hand sides of (3.4) and (3.5) agree. $\square$

In particular part ii) of the lemma asserts that for $\gamma \in \widehat{\Gamma}$,

$$\ln \left(\text{Iw}_{\eta(s)\widehat{A}}(\gamma \eta(s)g) \right) = \ln(\eta(s)^w) + \ln(\text{Iw}_{\widehat{A}}(g)^w) + \ln(\text{Iw}_{\widehat{A}}^{\mathbb{A}}(w u_w)) \quad (3.9)$$

for some element $u_w \in U_{w, \mathbb{A}}$, where the logarithm map $\ln$ was defined in (2.70) and (2.86).

3.3. We next establish some estimates on the individual terms in (3.9) for later use in section 4. For any parameter $t > 0$ let

$$\widehat{A}_t := \{h \in \widehat{A} \mid h^{a_i} > t \text{ for each } i = 1, \dots, \ell + 1\}, \quad (3.10)$$

and let $\widehat{U}_{\mathcal{D}}$ be a fundamental domain for the action of $\widehat{\Gamma} \cap \widehat{U}$ acting on $\widehat{U}$ by left translation. Siegel sets for $\eta(s)\widehat{G}$ were defined in [6] as sets of the form

$$\widehat{\mathfrak{S}}_t := \widehat{U}_{\mathcal{D}} \eta(s) \widehat{A}_t \widehat{K}, \quad (3.11)$$

for some choices of $t > 0$ and $\widehat{U}_{\mathcal{D}}$.

According to (2.42) and (2.43), any element $X \in \widehat{\mathfrak{h}}$ can be decomposed as

$$X = X_{cl} + \langle \Lambda_{\ell+1}, X \rangle h_t, \quad \text{with } X_{cl} \in \mathfrak{h} \text{ and } \langle \Lambda_{\ell+1}, X \rangle \in \mathbb{R}, \quad (3.12)$$

where we recall that $\Lambda_{\ell+1} : \widehat{\mathfrak{h}} \rightarrow \mathbb{R}$ from (2.18) is the $(\ell + 1)$ -st fundamental weight (it is trivial on all classical coroots). Note that because h_t and $\mathbf{c}$ are nonzero multiples of each other, properties (2.25) and (2.27) assert that all classical roots $\alpha \in \Delta$, classical coroots, fundamental weights $\omega_1, \dots, \omega_{\ell}$,

and ρ from (2.4) extend trivially to $\mathbb{R}h_t \oplus \mathbb{R}\mathbf{D}$. The classical component X_{cl} can be expanded as a linear combination

$$X_{cl} = \sum_{j=1}^{\ell} \langle \omega_j, X \rangle h_j \quad (3.13)$$

of the coroot basis $\{h_1, \dots, h_\ell\}$ of $\mathfrak{h}$ using (2.27). Because of property (2.5) we have that

$$\langle \rho, X \rangle = \langle \rho, X_{cl} \rangle = \sum_{j=1}^{\ell} \langle \omega_j, X \rangle. \quad (3.14)$$

With the above notation, we can now state the following result, which is one of the main technical tools in this paper.

Theorem. Fix an element $\eta(s)g$ of a Siegel set $\widehat{\mathfrak{S}}_t \subset \eta(s)\widehat{G}$ as in (3.11). Recall from (2.81) that $\eta(s) = \exp(r\mathbf{D})$ for $s = e^r$, $r \in \mathbb{R}_{>0}$. Then there exist positive constants $E = E(s, t, \Delta)$, $C_1 = C_1(s, t, \Delta)$, $C_2(s, t, \Delta)$ depending only on s , t , and the underlying classical root system Δ , and another positive constant $C_3 = C_3(g, s, t, \Delta)$ depending also on g with the following properties: for any element $w \in W^\theta$ as in (2.59) and $u_w \in U_{w, \mathbb{A}}$, the vectors H_1, H_2 , and $H_3 \in \widehat{\mathfrak{h}}$ defined by

$$H_1 := \ln(\text{Iw}_{\widehat{A}}^{\mathbb{A}}(wu_w)), \quad H_2 := \ln(\eta(s)^w) - r\mathbf{D}, \quad \text{and} \quad H_3 := \ln(\text{Iw}_{\widehat{A}}(g)^w) \quad (3.15)$$

satisfy the inequalities

$$\langle \omega_j, H_1 \rangle \geq 0 \quad \text{and} \quad \langle \omega_j, H_2 \rangle \geq 0 \quad \text{for } j = 1, \dots, \ell, \quad (3.16)$$

$$|\langle \omega_j, H_3 \rangle| \leq E \quad \text{for } j = 1, \dots, \ell, \quad (3.17)$$

and

$$\begin{aligned} \langle \Lambda_{\ell+1}, H_1 \rangle &\geq -C_1 (\ell(w) + 1) \langle \rho, H_1 \rangle, \\ \langle \Lambda_{\ell+1}, H_2 \rangle &\geq -C_2 (\ell(w) + 1) \langle \rho, H_2 \rangle, \\ \text{and} \quad \langle \Lambda_{\ell+1}, H_3 \rangle &\geq -C_3 (\ell(w) + 1). \end{aligned} \quad (3.18)$$

Remark: In fact, one can choose $C_1 = 1$ in the simply-laced case (as the proof below will show). The dependence of C_3 on g is relatively mild, and only enters through $\langle \Lambda_{\ell+1}, \ln(\text{Iw}_{\widehat{A}}(g)) \rangle$ in (3.49). In particular it is locally uniform in g .

3.4. The proof of the above inequalities will occupy the rest of this section. We shall first draw a corollary which will be useful in the sequel. Recall from (2.75) that the group $\widehat{A}$ has a direct product decomposition

$$\widehat{A} = \widehat{A}_{cen} \times \widehat{A}_{cl}, \quad (3.19)$$

where $\widehat{A}_{cen}$ is the connected component of the one-dimensional central torus of $\widehat{G}$ and $\widehat{A}_{cl}$ is the connected component of the split torus from the finite dimensional group G underlying $\widehat{G}$. As before we identify $\mathbb{R}_{>0}$ with $\widehat{A}_{cen}$ via the map $s \mapsto h_t(s)$. There is a natural projection from $\eta(s)\widehat{A}$ onto $\widehat{A}$, and subsequently onto each of the factors $\widehat{A}_{cen}$ and $\widehat{A}_{cl}$ in the decomposition (3.19).

Corollary. *There exist constants $C, D > 0$, depending only on s, t , the underlying classical root system Δ , and locally uniformly in g , with the following property: for any $w \in W^\theta$, $\gamma \in \widehat{\Gamma} \cap \widehat{B}w\widehat{B}$ and $\eta(s)g \in \widehat{\mathfrak{S}}_t$ we have the estimate*

$$x \geq Dy^{-(C(\ell(w)+1))^{-1}}, \quad (3.20)$$

where $a \in \widehat{A}_{cl}$ and $y \in \widehat{A}_{cen} \cong \mathbb{R}_{>0}$ are the projections of

$$\text{Iw}_{\eta(s)\widehat{A}}(\gamma\eta(s)g) \in \eta(s)\widehat{A} \quad (3.21)$$

onto $\widehat{A}_{cl}$ and $\widehat{A}_{cen}$, respectively, and $x = a^p \in \mathbb{R}_{>0}$.

Proof. We start by using part ii) of Lemma 3.2 (in particular, its consequence (3.9)) to write the logarithm of (3.21) as $rD + H_1 + H_2 + H_3$, where H_1, H_2 , and H_3 are defined in (3.15). This allows us to factor

$$x = x_1 x_2 x_3 \quad \text{and} \quad y = y_1 y_2 y_3, \quad (3.22)$$

where $x_i = e^{\langle \rho, H_i \rangle}$ and $y_i = e^{\langle \Lambda_{\ell+1}, H_i \rangle}$ for $i = 1, 2, 3$.

From the inequalities (3.18), there exists a constant $C > 0$ depending only on s, t, Δ , and g (in which it is locally uniform) such that

$$\begin{aligned} y_1 &\geq x_1^{-C(\ell(w)+1)}, \\ y_2 &\geq x_2^{-C(\ell(w)+1)}, \\ \text{and} \quad y_3 &\geq e^{-C(\ell(w)+1)}. \end{aligned} \quad (3.23)$$

Thus

$$y^{-(C(\ell(w)+1))^{-1}} = (y_1 y_2 y_3)^{-(C(\ell(w)+1))^{-1}} \leq x_1 x_2 e = x \frac{e}{x_3}. \quad (3.24)$$

Because of (2.4) and (3.17), the ratio $\frac{e}{x_3}$ is bounded above and below by positive constants depending only on s, t , and Δ , establishing (3.20). $\square$

3.5. In this subsection we prove the first inequalities of (3.16) and (3.18). We begin by recalling from [11, Lemma 6.1]³ that for any $w \in W^\theta$ and $u_w \in U_{w, \mathbb{A}}$,

$$H_1 = \ln(\text{Iw}_{\widehat{A}}^{\mathbb{A}}(wu_w)) = \sum_{\gamma \in \Delta_{-, w^{-1}}} c_\gamma h_\gamma \quad \text{with } c_\gamma \geq 0, \quad (3.25)$$

where $\Delta_{-, w^{-1}} = \{\gamma \in \widehat{\Delta}_- | w^{-1}\gamma > 0\}$ was introduced in (2.57). Recall from the definition (2.59) of W^θ that γ cannot be the negative of a simple classical root $\alpha_1, \dots, \alpha_\ell$, nor in their span. Since $\widehat{W}$ acts trivially $\widehat{\Delta}_I$, $\Delta_{-, w^{-1}}$ is a subset of $\widehat{\Delta}_W$ and its elements have the form (2.29).

Lemma. *Assume that $w \in W^\theta$ and $\gamma \in \Delta_{-, w^{-1}}$. Then γ has the form $\gamma = \beta + n\iota$ with $\beta \in \Delta_+$ and $n < 0$.*

³The equivalent result stated there uses the $\widehat{G} = \widehat{K}\widehat{A}\widehat{N}$ Iwasawa decomposition as opposed to the $\widehat{G} = \widehat{N}\widehat{A}\widehat{K}$ Iwasawa decomposition used here.

Proof. Since $\gamma < 0$ we must have that $n \leq 0$, and in fact $n < 0$ since it is not in the span of $\alpha_1, \dots, \alpha_\ell$. Let us write β as an integral linear combination $\beta = \sum_{i=1}^\ell d_i \alpha_i$ of the positive simple roots. If $d_1, \dots, d_\ell \leq 0$ then

$$w^{-1}\gamma = \sum_{i=1}^\ell d_i w^{-1}\alpha_i + n\iota \quad (3.26)$$

exhibits the positive root $w^{-1}\gamma$ as a nonpositive integral combination of positive roots, a contradiction. Thus at least one (and hence all) $d_i > 0$ and β is a positive root. $\square$

If we write $\gamma = \beta + n\iota \in \Delta_{-,w^{-1}}$ with $\beta \in \Delta_+$ and $n < 0$ as in the lemma, then (2.41) asserts the existence of a positive constant m such that $h_\gamma = h_\beta + mn h_\iota$. It follows from (3.25), (2.1), (2.27), (2.37-2.38), and the lemma that $\langle \omega_j, H_1 \rangle \geq 0$ for $j = 1, \dots, \ell$, which proves the first inequality of (3.16). Expand $H_1 = \sum_{j=1}^\ell \langle \omega_j, H_1 \rangle h_j + \langle \Lambda_{\ell+1}, H_1 \rangle h_\iota$ as in (3.12-3.13), so that

$$w^{-1}H_1 = \sum_{j=1}^\ell \langle \omega_j, H_1 \rangle h_{w^{-1}a_j} + \langle \Lambda_{\ell+1}, H_1 \rangle h_\iota \quad (3.27)$$

by (2.23). For each $j = 1, \dots, \ell$ we write the positive root $w^{-1}a_j$ as $\beta(j) + \kappa_j(w^{-1})\iota$, with $\beta(j) \in \Delta$ and $\kappa_j(w^{-1}) \geq 0$. Hence, again from (2.41) there exists a positive constant $m_{w^{-1}a_j}$ (which equals 1 in the simply-laced case, and which can only take on a finite number of values in general), such that

$$h_{w^{-1}a_j} = h_{\beta(j)} + m_{w^{-1}a_j} \kappa_j(w^{-1}) h_\iota. \quad (3.28)$$

The coefficient of h_ι in (3.27) is then given by the sum

$$\langle \Lambda_{\ell+1}, H_1 \rangle + \sum_{j=1}^\ell m_{w^{-1}a_j} \langle \omega_j, H_1 \rangle \kappa_j(w^{-1}). \quad (3.29)$$

On the other hand, from (2.47), (3.25), and the definition of $\gamma \in \Delta_{-,w^{-1}}$ it follows that $w^{-1}H_1$ is a nonnegative integral combination of positive coroots and so (3.29) is nonnegative, in particular

$$- \sum_{j=1}^\ell m_{w^{-1}a_j} \langle \omega_j, H_1 \rangle \kappa_j(w^{-1}) \leq \langle \Lambda_{\ell+1}, H_1 \rangle < 0. \quad (3.30)$$

The second inequality here comes from (3.25), the lemma, and (2.41). Lemma 2.9 asserts that $\kappa_j(w^{-1}) \leq \ell(w) + 1$. Since the values $m_{w^{-1}a_j}$ are bounded by an absolute constant (and again all equal to 1 in the simply-laced case), the first inequality of (3.18) now follows from (3.14).

3.6. Next, let us turn to the second of the inequalities listed in (3.16) and (3.18). Recall that we have defined H_2 in (3.15) through the formula

$$w(r\mathbf{D}) = r\mathbf{D} + H_2. \quad (3.31)$$

Now, let us write $w = \tilde{w}T_b$ with $\tilde{w} \in W$ and $b \in Q^\vee$. Decomposing $H_2 = H_{2,cl} + \langle \Lambda_{\ell+1}, H_2 \rangle h_\iota$ as in (3.12), formula (2.46) implies

$$\langle \Lambda_{\ell+1}, H_2 \rangle = -r \frac{(b,b)}{2} \quad \text{and} \quad H_{2,cl} = -r \tilde{w}b. \quad (3.32)$$

We claim that

$$d_j := \langle a_j, \tilde{w}b \rangle \leq 0 \quad \text{for } j = 1, \dots, \ell. \quad (3.33)$$

Indeed, from (2.45) we have that

$$w^{-1}a_j = T_{-b}\tilde{w}^{-1}a_j = \tilde{w}^{-1}a_j - \langle \tilde{w}^{-1}a_j, b \rangle \iota. \quad (3.34)$$

By definition (2.59), $w^{-1}a_j > 0$ for $w \in W^\theta$ and $j = 1, \dots, n$, hence

$$-\langle \tilde{w}^{-1}a_j, b \rangle \geq 0, \quad (3.35)$$

i.e., $d_j = \langle a_j, \tilde{w}b \rangle = \langle \tilde{w}^{-1}a_j, b \rangle \leq 0$ because of (2.21).

By Lemma 2.1 the inequality (3.33) implies that

$$q_j := \langle \omega_j, \tilde{w}b \rangle \leq 0 \quad \text{for } j = 1, \dots, \ell. \quad (3.36)$$

Hence,

$$\langle \omega_j, H_2 \rangle = \langle \omega_j, H_{2,cl} \rangle = -r \langle \omega_j, \tilde{w}b \rangle \geq 0 \quad \text{for } j = 1, \dots, \ell. \quad (3.37)$$

This proves the second inequality in (3.16).

Let us now turn to the second inequality of (3.18). From (3.33) and (3.36) we have the simultaneous expressions

$$\begin{aligned} \tilde{w}b &= \sum_{j=1}^{\ell} q_j h_j \quad \text{with } q_j \leq 0 \\ \text{and} \quad \tilde{w}b &= \sum_{j=1}^{\ell} d_j \omega_j^\vee \quad \text{with } d_j \leq 0, \end{aligned} \quad (3.38)$$

where ω_j and $\omega_j^\vee$ are defined in (2.1)-(2.2). Using (2.3) we find that

$$(\tilde{w}b, \tilde{w}b) = \sum_{i=1}^{\ell} \frac{2}{(\alpha_i, \alpha_i)} d_i q_i. \quad (3.39)$$

Since the denominators take on only a finite number of positive values, there exists a constant $B > 0$ such that

$$(\tilde{w}b, \tilde{w}b) \leq B \sum_{i=1}^{\ell} d_i q_i \leq -B \cdot \max_i |d_i| \cdot \sum_{j=1}^{\ell} q_j. \quad (3.40)$$

Applying Lemma 2.7 results in the estimate $\max_i |d_i| \leq C'(\ell(w) + 1)$ for some constant $C' > 0$ independent of w . Thus there exists some constant $C'' > 0$, again independent of w , such that

$$-\langle \Lambda_{\ell+1}, H_2 \rangle = r \frac{(b, b)}{2} = r \frac{(\tilde{w}b, \tilde{w}b)}{2} \leq -C'' (\ell(w) + 1) \sum_{j=1}^{\ell} r q_j. \quad (3.41)$$

On the other hand, we have from (2.5), (3.32), and (3.38) that

$$\langle \rho, H_{2,cl} \rangle = -\langle \rho, r\tilde{w}b \rangle = -r\langle \rho, \sum_{j=1}^{\ell} q_j h_j \rangle = -\sum_{j=1}^{\ell} r q_j. \quad (3.42)$$

The second inequality of (3.18) follows from (3.41) and (3.42).

3.7. Finally, we turn to (3.17) and the last inequality (3.18) in Theorem 3.3. Let $H_g := \ln(\text{Iw}_{\hat{A}}(g))$, and write

$$H_g = H_{g,cl} + \langle \Lambda_{\ell+1}, H_g \rangle h_l, \quad (3.43)$$

where $H_{g,cl} \in \mathfrak{h}$ as in (3.12). If we write $w = \tilde{w}T_b$ with $\tilde{w} \in W$ and $b \in Q^\vee$, we have

$$\begin{aligned} H_3 &:= wH_g = \tilde{w}T_b H_{g,cl} + \langle \Lambda_{\ell+1}, H_g \rangle h_l \\ &= \tilde{w}H_{g,cl} + (\langle \Lambda_{\ell+1}, H_g \rangle + (H_{g,cl}, b)) h_l, \end{aligned} \quad (3.44)$$

where we have used (2.46) and the fact that w fixes h_l . This implies

$$(H_3)_{cl} = \tilde{w}H_{g,cl}, \quad (3.45)$$

in particular.

Recall that we have assumed that $\eta(s)g \in \hat{\mathfrak{S}}_t$ with $t > 0$. Because of (3.10) we must have that

$$\text{Iw}_{\hat{A}}(\eta(s)g)^{a_i} = e^{\langle a_i, r\mathbf{D} + H_g \rangle} > t \quad \text{for } i = 1, 2, \dots, \ell + 1, \quad (3.46)$$

i.e., $\langle a_i, r\mathbf{D} + H_g \rangle > \ln(t)$ for $i = 1, \dots, \ell + 1$. Since (2.25)-(2.26) imply that $\langle a_i, h \rangle = 0$ for $h \in \mathbb{R}h_l \oplus \mathbb{R}\mathbf{D}$ and $i = 1, \dots, \ell$, we furthermore have that

$$\langle a_i, r\mathbf{D} + H_g \rangle = \langle a_i, H_{g,cl} \rangle > \ln(t) \quad \text{for } i = 1, \dots, \ell. \quad (3.47)$$

On the other hand, from (2.28) and (2.31) we also have that

$$\langle a_{\ell+1}, r\mathbf{D} + H_g \rangle = \langle -\alpha_0 + \mathfrak{t}, r\mathbf{D} + H_g \rangle = -\langle \alpha_0, H_{g,cl} \rangle + r > \ln(t), \quad (3.48)$$

where we recall that α_0 is the highest root for the finite-dimensional Lie algebra $\mathfrak{g}$, and as such is a positive linear combination of the roots a_i , $i = 1, \dots, \ell$. Thus, from (3.47) and (3.48) we conclude that as $\eta(s)g$ varies over the Siegel set $\hat{\mathfrak{S}}_t$, the quantity $|\langle a_i, H_{g,cl} \rangle|$ is bounded for $i = 1, \dots, \ell$ by a constant which depends only on t, r , and the root system Δ . Since $\tilde{w}$ varies over a finite set, inequality (3.17) now follows from (3.45) and Lemma 2.1.

Formula (3.44) also implies that

$$\langle \Lambda_{\ell+1}, H_3 \rangle = \langle \Lambda_{\ell+1}, H_g \rangle + (H_{g,cl}, b). \quad (3.49)$$

Using (3.17), Lemma 2.7, and the Cauchy-Schwartz inequality, we see that there exists a constant $C = C(s, t, \Delta)$ depending on such that $|(H_{3,cl}, b)| \leq C(\ell(w) + 1)$. Since $\langle \Lambda_{\ell+1}, H_g \rangle$ depends locally uniformly on H_g , the last inequality of (3.18) follows.

4 Entire absolute convergence on loop groups

In this section, we combine our analysis from section 3 with a decay estimate on cusp forms to conclude that cuspidal loop Eisenstein series are entire. We begin with some discussion of the structure of parabolic subgroups of loop groups.

4.1. Let k be any field and consider the group $\widehat{G}_k$ constructed in section 2.10. For any subset $\theta \subset \{1, \dots, \ell + 1\}$ let $\widehat{W}_\theta$ denote the subgroup of $\widehat{W}$ generated by the reflections $\{w_i | i \in \theta\}$, and define the parabolic subgroup $\widehat{P}_{\theta,k} \subset \widehat{G}_k$ as

$$\widehat{P}_{\theta,k} := \widehat{B}_k \widehat{W}_\theta \widehat{B}_k, \quad (4.1)$$

where each $w \in \widehat{W}_\theta$ is identified with a representative in $N(\widehat{H}_k)$ as in section 2.11. Thus $\widehat{P}_{\theta,k} = \widehat{G}_k$ when $\theta = \{1, \dots, \ell + 1\}$ (cf. (2.78)). Denote by $\widehat{U}_{\theta,k}$ the pro-unipotent radical of $\widehat{P}_{\theta,k}$. If $\theta \subsetneq \{1, \dots, \ell + 1\}$ the group $\widehat{W}_\theta$ is finite, and the group $\widehat{U}_{\theta,k}$ can be written as the intersection of $\widehat{U}_k$ with a conjugate by the longest element in $\widehat{W}_\theta$. Recall the subgroup $\widehat{H}_k \subset \widehat{G}_k$ defined at the beginning of section 2.11 as the diagonal operators with respect to the coherent basis $\mathcal{B}$. We now set

$$\begin{aligned} \widehat{H}_{\theta,k} &= \langle h \in \widehat{H}_k | h^{a_i} = 1 \text{ for all } i \in \theta \rangle \\ \text{and } H(\theta)_k &= \langle h_i(s) | \text{ for } i \in \theta \text{ and } s \in k^* \rangle. \end{aligned} \quad (4.2)$$

Analogously to (2.75), there is an almost direct product decomposition $\widehat{H}_k = \widehat{H}_{\theta,k} \times H(\theta)_k$. Let

$$L_{\theta,k} = \langle \chi_\beta(s) | \beta \in [\theta], s \in k \rangle, \quad (4.3)$$

where $[\theta]$ denotes the set of all roots in $\widehat{\Delta}$ which can be expressed as linear combinations of elements of θ . We now define

$$M_{\theta,k} = L_{\theta,k} \widehat{H}_{\theta,k}, \quad (4.4)$$

its semi-simple quotient

$$L'_{\theta,k} = M_{\theta,k} / Z(M_{\theta,k}) = L_{\theta,k} / (Z(M_{\theta,k}) \cap L_{\theta,k}), \quad (4.5)$$

and the natural projection

$$\pi_{L'_{\theta,k}} : M_{\theta,k} \rightarrow L'_{\theta,k}. \quad (4.6)$$

With the notation above, $\widehat{P}_{\theta,k}$ decomposes into the semidirect product

$$\widehat{P}_{\theta,k} = \widehat{U}_{\theta,k} \rtimes M_{\theta,k} \quad (4.7)$$

(see [7, Theorem 6.1]).

4.2. We now suppose that $k = \mathbb{R}$ and continue our convention of dropping the subscript k when referring to real groups. Recall the subgroup $\widehat{A}$ of $\widehat{H}$ defined in section 2.10. Setting $\widehat{A}_\theta = \widehat{H}_\theta \cap \widehat{A}$ and $A(\theta) = H(\theta) \cap \widehat{A}$, it has the direct product decomposition

$$\widehat{A} = \widehat{A}_\theta \times A(\theta) \quad (4.8)$$

as a consequence of the fact the Cartan submatrix corresponding to θ is positive definite [22, Lemma 4.4].

The decomposition (4.4) is not a direct product, but can be refined to one as follows. Define

$$M_\theta^1 = \bigcap_{\chi \in X(M_\theta)} \ker(\chi^2), \quad (4.9)$$

where $X(M_\theta)$ denotes the set of real algebraic characters of M_θ .⁴ Then the direct product decomposition

$$M_\theta = M_\theta^1 \times \widehat{A}_\theta. \quad (4.10)$$

holds.

The group $\widehat{G}$ has an Iwasawa decomposition with respect to the parabolic $\widehat{P}_\theta$,

$$\widehat{G} = \widehat{P}_\theta \widehat{K} = \widehat{U}_\theta M_\theta \widehat{K}. \quad (4.11)$$

In contrast to (2.80), this Iwasawa decomposition is typically not unique since $K_\theta := \widehat{K} \cap M_\theta$ may be nontrivial; in general only the map

$$\text{Iw}_{M_\theta} : \widehat{G} \rightarrow M_\theta / K_\theta \quad (4.12)$$

is well-defined. Noting that K_θ is a compact subgroup of the finite-dimensional group M_θ , one in fact has that $K_\theta \subset M_\theta^1$. This and the direct product (4.10) allow us to define the map

$$\text{Iw}_{\widehat{A}_\theta} : \widehat{G} \rightarrow M_\theta / M_\theta^1 \cong \widehat{A}_\theta, \quad (4.13)$$

which factors through Iw_{M_θ} .

4.3. The finite-dimensional, real semi-simple Lie group L'_θ admits an Iwasawa decomposition

$$L'_\theta = U'_\theta A'(\theta) K'_\theta \quad (4.14)$$

of its own, where $U'_\theta \subset L'_\theta$ is a unipotent subgroup, $A'(\theta) = \pi_{L'_\theta}(A(\theta))$, and

$$K'_\theta = \pi_{L'_\theta}(K_\theta) = \pi_{L'_\theta}(\widehat{K} \cap M_\theta). \quad (4.15)$$

The projection from L'_θ onto $A'(\theta)$ will be denoted by

$$\text{Iw}_{A'(\theta)} : L'_\theta \rightarrow A'(\theta). \quad (4.16)$$

The factors $\widehat{A}_\theta$ and $A(\theta)$ have trivial intersection in (4.8) and the factors in (4.10) commute. Therefore the intersection $Z(M_\theta) \cap A(\theta) = Z(M_\theta^1) \cap \widehat{A}_\theta \cap A(\theta)$ is also trivial, and hence the map $\pi_{L'_\theta}$ induces an isomorphism

$$\pi_{L'_\theta} : A(\theta) \cong A'(\theta). \quad (4.17)$$

⁴The group M_θ^1 was denoted $\widetilde{L}_\theta$ in [14]

Define the map $\text{Iw}_{A(\theta)}$ as the composition of the Iwasawa $\widehat{A}$ -projection of $\widehat{G}$ defined after (2.80) together with the projection onto the second factor in (4.8),

$$\text{Iw}_{A(\theta)} : \widehat{G} \xrightarrow{\text{Iw}_{\widehat{A}}} \widehat{A} \rightarrow A(\theta). \quad (4.18)$$

Then the composition

$$\widehat{G} \xrightarrow{\text{Iw}_{M_\theta}} M_\theta/K_\theta \xrightarrow{\pi_{L'_\theta}} L'_\theta/K'_\theta \xrightarrow{\text{Iw}_{A'(\theta)}} A'(\theta) \quad (4.19)$$

coincides with $\pi_{L'_\theta} \circ \text{Iw}_{A(\theta)}$.

4.4. We shall fix the choice $\theta = \{1, \dots, \ell\}$ for the remainder of this section, so that $\widehat{W}_\theta = W$. The groups $\widehat{A}_\theta$ and $A(\theta)$ are then respectively isomorphic to the groups $\widehat{A}_{cen}$ and $\widehat{A}_{cl}$ defined just after (2.74). We will thus denote the maps $\text{Iw}_{\widehat{A}_\theta}$ and $\text{Iw}_{A(\theta)}$ simply by $\text{Iw}_{\widehat{A}_{cen}}$ and $\text{Iw}_{\widehat{A}_{cl}}$, respectively. Since $\eta(s)$ normalizes $\widehat{U}$ we may extend $\text{Iw}_{\widehat{A}_{cen}}$ and $\text{Iw}_{\widehat{A}_{cl}}$ to maps on the slice $\eta(s)\widehat{G}$ which we continue to denote by the same names,

$$\text{Iw}_{\widehat{A}_{cen}} : \eta(s)\widehat{G} \rightarrow \widehat{A}_{cen} \quad \text{and} \quad \text{Iw}_{\widehat{A}_{cl}} : \eta(s)\widehat{G} \rightarrow \widehat{A}_{cl}, \quad (4.20)$$

similarly to (2.84). Likewise we may furthermore define the map

$$\text{Iw}_{M_\theta} : \eta(s)\widehat{G} \rightarrow M_\theta/K_\theta = M_\theta/(\widehat{K} \cap M_\theta), \quad (4.21)$$

noting from (2.82) that $\eta(s)$ acts trivially on $M_\theta = M_{\{1, \dots, \ell\}}$.

Recall that the group $\widehat{A}_{cen}$ consists of the elements $h_i(s)$ for $s > 0$. For a complex number $\nu \in \mathbb{C}$ define $h_i(s)^\nu = s^\nu$, and consider the function

$$\begin{aligned} \eta(s)\widehat{G} &\rightarrow \mathbb{C}^* \\ \eta(s)g &\mapsto \text{Iw}_{\widehat{A}_{cen}}(\eta(s)g)^\nu. \end{aligned} \quad (4.22)$$

Let $\widehat{P}$ be the parabolic $\widehat{P}_\theta = \widehat{P}_{\{1, \dots, \ell\}}$. The following convergence theorem for Eisenstein series⁵ has been proven by the first named author.

Theorem (Theorem 3.2 in [14]). *The Eisenstein series*

$$\sum_{\gamma \in (\widehat{\Gamma} \cap \widehat{P}) \backslash \widehat{\Gamma}} \text{Iw}_{\widehat{A}_{cen}}(\gamma\eta(s)g)^\nu \quad (4.23)$$

converges absolutely for $\text{Re } \nu > 2h^\vee$, where $h^\vee = \langle \rho, h_i \rangle$ is the dual Coxeter number. Moreover, the convergence is uniform when g is constrained to a subset of the form $\widehat{U}_\mathscr{D} \widehat{A}_{cpt} \widehat{K}$, where $\widehat{A}_{cpt} \subset \widehat{A}$ is compact and $\widehat{U}_\mathscr{D}$ is as in (3.11).

4.5. Our aim is to prove the convergence of the cuspidal analogs of (4.23) for all $\nu \in \mathbb{C}$. These were defined in [14], where they were shown to converge in a right half plane. For any right K_θ -invariant function $f : M_\theta \rightarrow \mathbb{C}$ the assignment

$$\eta(s)g \mapsto f(\text{Iw}_{M_\theta}(\eta(s)g)) \quad (\text{cf. (4.21)}) \quad (4.24)$$

⁵The result in [14] states absolute convergence for the left half plane $\text{Re } (\nu) < -2h^\vee$, since the order of the Iwasawa decomposition there is reversed.

is a well-defined function from $\eta(s)\widehat{G}$ to $\mathbb{C}$. Recall that the finite-dimensional semisimple group L'_θ was defined in (4.5) together with the projection map $\pi_{L'_\theta} : M_\theta \rightarrow L'_\theta$. Set

$$\Gamma_\theta := \widehat{\Gamma} \cap M_\theta \quad (4.25)$$

and $\Gamma'_\theta = \pi_{L'_\theta}(\Gamma_\theta)$. Recall that $K'_\theta = \pi_{L'_\theta}(K_\theta)$ from (4.15) and let ϕ be a K'_θ -invariant cusp form on $\Gamma'_\theta \backslash L'_\theta$. Set

$$\Phi = \phi \circ \pi_{L'_\theta} : \Gamma_\theta \backslash M_\theta \rightarrow \mathbb{C}, \quad (4.26)$$

which is right K_θ -invariant. The *cuspidal loop Eisenstein series* from [14] is defined as the sum

$$E_{\phi, \nu}(\eta(s)g) := \sum_{\gamma \in (\widehat{\Gamma} \cap \widehat{P}) \backslash \widehat{\Gamma}} \text{Iw}_{\widehat{A}_{\text{cen}}}(\gamma \eta(s)g)^\nu \Phi(\text{Iw}_{M_\theta}(\gamma \eta(s)g)). \quad (4.27)$$

We can now state the main result of this paper

Theorem. *The cuspidal loop Eisenstein series, $E_{\phi, \nu}(\eta(s)g)$ converges absolutely for any $\nu \in \mathbb{C}$. Moreover, the convergence is uniform when g is constrained to a subset of the form $\widehat{U}_\mathscr{D} \widehat{A}_{\text{cpt}} \widehat{K}$, where $\widehat{A}_{\text{cpt}} \subset \widehat{A}$ is compact and $\widehat{U}_\mathscr{D}$ is as in (3.11).*

4.6. The proof of Theorem 4.5 is based on two main ingredients, namely the inequalities proved in Theorem 3.3 and also the following decay estimate. To state it, we keep the notation as in §2.1 – 2.3. Let $\phi : \Gamma \backslash G \rightarrow \mathbb{C}$ be a K -finite cusp form. The following result can be deduced from Theorem A.3.1 in the appendix. Stronger results are known for K -fixed cusp forms due to Bernstein and Krötz-Opdam [23], and those are in fact sufficient for the applications here since we only consider Eisenstein series induced from K -fixed cusp forms. Their techniques extend to the K -finite setting, but have not been published. As mentioned in the introduction, we anticipate the K -finite statement will be useful for proving the convergence of K -finite loop Eisenstein series once a definition has been given, and so we give a complete proof of the following result in the appendix (using a different argument).

Theorem. *Let $\phi \in \Gamma \backslash G$ be a K -finite cusp form. Then there exists a constant $C > 0$ which depends only on G and ϕ such that for every natural number $N \geq 1$, we have*

$$\phi(g) \leq (CN)^{CN} \text{Iw}_A(g)^{-N\rho}. \quad (4.28)$$

This is exactly the statement of (A.26) if g is in some Siegel set $\mathfrak{S}_t$. In general, if $g \in G$ we can find $\gamma \in \Gamma$ such that $\gamma g \in \mathfrak{S}_t$. By Remark 2.3.1 we have that

$$\text{Iw}_A(\gamma g)^\rho \geq \text{Iw}_A(g)^\rho, \quad (4.29)$$

and so the asserted estimate for $\phi(g) = \phi(\gamma g)$ reduces to the estimate on $\mathfrak{S}_t$.

4.6.1. Remark The explicit N -dependence in the estimate (4.28) is needed in showing the boundedness of (4.40) below. Indeed, since N there depends on the Bruhat cell, the classical rapid decay statements (in which $(CN)^{CN}$ is replaced by some constant depending on N) are insufficient.

4.7. This subsection contains the proof of Theorem 4.5. Since cusp forms are bounded, the convergence for $\text{Re } \nu > 2h^\vee$ follows from that of (4.23) (as observed by Garland in [14]). We shall

prove the theorem for $\operatorname{Re} \nu \leq 2h^\vee$ by leveraging the decay of the cusp form to dominate the series (4.27) by a convergent series of the form (4.23), but with ν replaced by some $\nu_0 \gg \operatorname{Re} \nu$.

4.7.1. Step 1: Main Analytic Estimate. For $w \in W^\theta$ let $\widehat{\Gamma}(w) = \widehat{\Gamma} \cap \widehat{P}w\widehat{P}$, which is independent of the representative in $N(\widehat{H})$ taken for w . Each set $\widehat{\Gamma}(w)$ is invariant under $\widehat{\Gamma} \cap \widehat{P}$. Using the Bruhat decomposition, the Eisenstein series can then be written as

$$E_{\phi, \nu}(\eta(s)g) = \sum_{w \in W^\theta} \sum_{\gamma \in (\widehat{\Gamma} \cap \widehat{P}) \backslash \widehat{\Gamma}(w)} \operatorname{Iw}_{\widehat{A}_{cen}}(\gamma\eta(s)g)^\nu \Phi(\operatorname{Iw}_{M_\theta}(\gamma\eta(s)g)). \quad (4.30)$$

Recall that we have assumed $\operatorname{Re} \nu \leq 2h^\vee$. Choose $\nu_0 \in \mathbb{R}$ with $\nu_0 > 2h^\vee \geq \operatorname{Re} \nu$, so that the series (4.23) converges when ν is replaced by ν_0 . Consider the inner sum on the right hand side of (4.30) for a fixed element $w \in W^\theta$,

$$\sum_{\gamma \in (\widehat{\Gamma} \cap \widehat{P}) \backslash \widehat{\Gamma}(w)} \operatorname{Iw}_{\widehat{A}_{cen}}(\gamma\eta(s)g)^\nu \Phi(\operatorname{Iw}_{M_\theta}(\gamma\eta(s)g)). \quad (4.31)$$

In the notation of Corollary 3.4, let

$$y = \operatorname{Iw}_{\widehat{A}_{cen}}(\gamma\eta(s)g) \in \mathbb{R}_{>0} \quad \text{and} \quad x = \operatorname{Iw}_{\widehat{A}_{cl}}(\gamma\eta(s)g)^\rho \in \mathbb{R}_{>0}. \quad (4.32)$$

By the definition of Φ given in (4.26), $\Phi(\operatorname{Iw}_{M_\theta}(\gamma\eta(s)g)) = \phi(\pi_{L'_\theta}(\operatorname{Iw}_{M_\theta}(\gamma\eta(s)g)))$. Since $\widehat{A}_{cl} = A(\theta)$, we may conclude from what we have noted after (4.18) that

$$\operatorname{Iw}_{A'(\theta)}(\pi_{L'_\theta}(\operatorname{Iw}_{M_\theta}(\gamma\eta(s)g))) = \pi_{L'_\theta}(\operatorname{Iw}_{\widehat{A}_{cl}}(\gamma\eta(s)g)). \quad (4.33)$$

Applying Theorem 4.6 we conclude that there exists a constant $C_1 > 0$ such that

$$\Phi(\operatorname{Iw}_{M_\theta}(\gamma\eta(s)g)) = \phi(\pi_{L'_\theta}(\operatorname{Iw}_{M_\theta}(\gamma\eta(s)g))) \leq (C_1 N)^{C_1 N} x^{-N} \quad (4.34)$$

for any $N \in \mathbb{Z}_{>0}$.

4.7.2. Step 2: Comparing the central contribution. From Corollary 3.4 there exist constants $C_2, D > 0$ independent of $\gamma \in \widehat{\Gamma} \cap \widehat{B}w\widehat{B}$, but depending locally uniformly on g , such that

$$x \geq D y^{-(C_2(\ell(w)+1))^{-1}}. \quad (4.35)$$

Choose a positive real d such that $dC_2 \in \mathbb{Z}_{>0}$ and

$$d > \nu_0 - \operatorname{Re} \nu > 0, \quad (4.36)$$

and let $N = dC_2(\ell(w) + 1)$. From (4.35) we obtain

$$x^{-N} y^{\operatorname{Re} \nu - \nu_0} \leq D^{-N} y^{\frac{N}{C_2(\ell(w)+1)}} y^{\operatorname{Re} \nu - \nu_0} = D^{-N} y^d y^{\operatorname{Re} \nu - \nu_0}. \quad (4.37)$$

Thus

$$\begin{aligned} \operatorname{Iw}_{\widehat{A}_{cen}}(\gamma\eta(s)g)^{\operatorname{Re} \nu} \Phi(\operatorname{Iw}_{M_\theta}(\gamma\eta(s)g)) &= y^{\operatorname{Re} \nu} \Phi(\operatorname{Iw}_{M_\theta}(\gamma\eta(s)g)) \\ &\leq (C_1 N)^{C_1 N} y^{\operatorname{Re} \nu} x^{-N} \\ &\leq (C_1 N)^{C_1 N} y^{\nu_0} (x^{-N} y^{\operatorname{Re} \nu - \nu_0}) \\ &\leq (C_1 N)^{C_1 N} y^{\nu_0} D^{-N} y^d, \end{aligned} \quad (4.38)$$

where $a = d + \operatorname{Re} v - v_0$ is positive by (4.36).

On the other hand,

$$\ln y = \langle \Lambda_{\ell+1}, \ln(\gamma\eta(s)g) \rangle = \langle \Lambda_{\ell+1}, H_1 \rangle + \langle \Lambda_{\ell+1}, H_2 \rangle + \langle \Lambda_{\ell+1}, H_3 \rangle \quad (4.39)$$

in the notation of Theorem 3.3. The second inequality in (3.30) states that $\langle \Lambda_{\ell+1}, H_1 \rangle < 0$. From (3.32) and Lemma 2.7, the term $\langle \Lambda_{\ell+1}, H_2 \rangle$ is bounded above by a quadratic polynomial in $\ell(w)$ with strictly negative quadratic term. Furthermore, for fixed $\eta(s)g \in \widehat{\mathfrak{S}}_t$, it follows from (3.49) that $|\langle \Lambda_{\ell+1}, H_3 \rangle|$ is bounded above by a linear polynomial in $\ell(w)$ with coefficients that depend locally uniformly in g . Hence, in total y is bounded above by an expression of the form $e^{p(\ell(w))}$ where $p(\cdot)$ is a quadratic polynomial with strictly negative quadratic term. By our choice of N the term $D^{-N}(C_1N)^{C_1N}$ is bounded above by a constant times $e^{c'\ell(w)\ln \ell(w)}$ for some positive constant c' . Hence, the expression

$$(C_1N)^{C_1N} D^{-N} y^a \quad (4.40)$$

is bounded as $\ell(w) \rightarrow \infty$, and so we can (4.27) by a sum of the form

$$\sum_{w \in W^\theta} \sum_{\gamma \in (\widehat{\Gamma} \cap \widehat{P}) \setminus \widehat{\Gamma}(w)} \operatorname{Iw}_{\widehat{A}_{cen}}(\gamma\eta(s)g)^{v_0} = \sum_{\gamma \in (\widehat{\Gamma} \cap \widehat{P}) \setminus \widehat{\Gamma}} \operatorname{Iw}_{\widehat{A}_{cen}}(\gamma\eta(s)g)^{v_0}. \quad (4.41)$$

The convergence of (4.27) then follows from the fact that $v_0 > 2h^\vee$ is in the range of convergence of (4.23).

4.8. Instead of working over $\mathbb{Q}$, we can also study cuspidal loop Eisenstein series over a function field F of a smooth projective curve X over a finite field. It was in this setting that Braverman and Kazhdan originally noticed the entirety of cuspidal Eisenstein series [4]. In fact, they observed something stronger which is peculiar to the function field setting. Namely, for every fixed element in the appropriate symmetric space, the Eisenstein series is a finite sum. Their argument is geometric in nature and relies on estimating the number of points in certain moduli spaces which arise in a geometric construction of Eisenstein series.

Alternatively, one can also easily adapt the arguments in this paper to the function field setting to reprove the result of Braverman and Kazhdan. We briefly indicate the argument here. As in the proof of Theorem 4.5, we first break up the Eisenstein series into a sum over Bruhat cells indexed by $w \in W^\theta$. On each such cell, it is easy to see using the arguments in [11, Lemma 2.5] that there can be at most finitely many elements whose classical Iwasawa component lies in the support of any compactly supported function on $(Z(M_{\theta, \mathbb{A}_F})M_{\theta, F}) \backslash M_{\theta, \mathbb{A}_F} / (K_{\mathbb{A}_F} \cap M_{\theta, \mathbb{A}_F})$. It is a result of Harder (see [27, Lemma I.2.7]) that cusp forms do indeed have compact support on their fundamental domain. Hence, only finitely many terms contribute to the cuspidal loop Eisenstein series on each cell. Moreover, one can use an analogue of our main Iwasawa inequalities Theorem 3.3 to show that cells corresponding to w with $\ell(w)$ sufficiently large cannot contribute *any* non-zero elements to the Eisenstein series. Indeed, as in the proof of Theorem 4.5, one may again show that as $\ell(w) \rightarrow \infty$ the central term of the Iwasawa component is bounded above by $e^{p(\ell(w))}$ where $p(\cdot)$ is a quadratic polynomial with strictly negative quadratic term. On the other hand, the analogue of (4.35) dictates that as $\ell(w) \rightarrow \infty$ the central piece grows smaller and the classical piece must grow large and therefore eventually lie outside the support of the cusp form.

A K -finite cusp forms decay exponentially

A.1 Introduction and statement of main result

In this appendix we give a proof of Theorem 4.6. This rapid decay estimate with uniform constants will be deduced from an exponential-type decay statement. Historically rapid decay estimates (without uniformity) were established in order to circumvent the apparent difficulty of proving the exponential decay estimates familiar in the classical theory of automorphic forms for $SL(2, \mathbb{R})$. As our argument here indicates, one gets qualitatively similar estimates for K -finite cusp forms on general groups by injecting a small amount of one-variable hard analysis. Stronger exponential decay results were proven by Bernstein using the more sophisticated technique of holomorphic continuation of representations (for which we refer to Krötz-Opdam [23] for K -fixed cusp forms, and unpublished notes of theirs for K -finite forms). Although we state and prove the results here in the setting of Chevalley groups used in the body of the paper, the results and the methods are applicable to general groups.

The application to Eisenstein series in the body of this paper uses only K -fixed forms, since (as noted in the introduction) at present it is difficult to formulate a definition of Eisenstein series induced from general K -finite forms without further developments in the representation theory of affine loop groups. However, this issue seems independent of the analysis we used to demonstrate convergence, and so the bounds we prove here should ultimately play the same role in showing the everywhere absolute convergence (and hence entirety) of those series as well. Since this estimate does not appear in the literature we have chosen to include this appendix.

Let G be a Chevalley group over $\mathbb{R}$ and Γ an arithmetic subgroup with respect to the Chevalley structure. Other notations used here will have the same meaning as in the main body of this paper (see §2A, in particular (2.8)).

Theorem. *Let ϕ be a K -finite cusp form on $\Gamma \backslash G$. For any fixed $t > 0$ there exist constants $c, d > 0$ such that ϕ satisfies the estimate*

$$\phi(ak) \leq \exp\left(-c \left|\max_{\alpha \in \Delta_+} a^\alpha\right|^d\right) \quad \text{for all } a \in A_t. \quad (\text{A.1})$$

The constant $c > 0$ is certainly necessary, as can already be seen in the context of classical modular forms on the complex upper half plane. Indeed, a cusp form invariant under parabolic transformations $z \mapsto z + w$ is bounded by a multiple of $e^{-2\pi w^{-1} \text{Im}(z)}$. Bernstein's holomorphic continuation method mentioned above shows that $d \geq 1$, in consonance with classical modular forms. Up to logarithmic factors, we are not aware of any examples where d is not 1, though we make some comments in §A.6 about where they might occur. Even with $d \neq 1$ the estimate (A.1) could be regarded as exponential decay from the vantage point of the Lie algebra $\mathfrak{h}$ of A , where dilation by a constant multiple has the effect of scaling d (hence the title of this appendix).

The rest of the appendix is arranged as follows: in §A.2 we first study the decay of a certain function on G whose properties will be crucial later in the argument. Combining this estimate with a slight enhancement of a result of Harish-Chandra (proposition A.2.4), we then retrace our way in §A.3 through the usual proof of rapid decay to obtain the asserted rapid decay with uniform constants. We then deduce exponential decay from this estimate in §A.4. Finally, in §A.5-A.6 we aim to place this result in a broader context, by giving an example which demonstrates the necessity of the K -finite condition and explaining the importance of estimates of the form (A.1) in string theory.

A.2 Derivative estimates on a particular function

A.2.1. We begin by estimating the norm of a certain function on the real line. Let $\sigma \in C_c^\infty(\mathbb{R})$ be the “bump” function

$$\sigma(x) := \begin{cases} e^{-1/(1-x^2)}, & |x| < 1, \\ 0, & \text{otherwise,} \end{cases} \quad (\text{A.2})$$

which is supported in the interval $[-1, 1]$.

Lemma. *The function σ satisfies the L^1 -norm estimate*

$$\left\| \frac{d^N \sigma}{dx^N} \right\|_1 \ll (2N)^{2N} \quad (\text{A.3})$$

for positive integers N .

Proof. A standard application of the saddle point method⁶ shows that its Fourier transform satisfies the bound $\widehat{\sigma}(r) \ll r^{-3/4} e^{-\sqrt{2\pi}r}$. In particular

$$\begin{aligned} \|\widehat{\sigma}(r)r^N\|_2^2 &= \int_{\mathbb{R}} \widehat{\sigma}(r)^2 r^{2N} dr \ll \int_0^\infty e^{-\sqrt{8\pi}r} r^{2N-3/2} dr \\ &\ll (2\pi)^{-2N} \Gamma(4N-1) \leq (2\pi)^{-2N} (4N)! \ll (2\pi)^{-2N} (2N)^{4N} \end{aligned} \quad (\text{A.4})$$

and so by Cauchy-Schwartz, the compact support of σ , and Parseval’s Theorem we have

$$\left\| \frac{d^N \sigma}{dx^N} \right\|_1 \ll \left\| \frac{d^N \sigma}{dx^N} \right\|_2 = (2\pi)^N \|\widehat{\sigma}(r)r^N\|_2 \ll (2N)^{2N}. \quad (\text{A.5})$$

□

For further reference we note that

$$\left\| \frac{d^N \sigma}{dx^N} \sigma(c^{-1}x) \right\|_1 \ll (2Nc^{-1})^{2Nc^{-1}} \quad (\text{A.6})$$

for $c < 1$. Incidentally, one might wonder whether replacing σ in (A.2) with a different choice of function might lead to significantly better estimates later on. Other than improving the constant “2” in (A.3) this is impossible [32], consistent with the fact one cannot improve upon the known exponential decay of classical holomorphic cusp forms.

A.2.2. We next use σ construct a family of bump functions on our group G with Lie algebra $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$. Let $\{Y_1, Y_2, \dots, Y_k\}$, $\{H_1, \dots, H_r\}$, and $\{X_1, X_2, \dots, X_k\}$ denote bases of these three summands, respectively. Combined they give a basis

$$\mathcal{B} = \{Y_1, \dots, Y_k, H_1, \dots, H_r, X_1, \dots, X_k\} \quad (\text{A.7})$$

of the $d = 2k + r$ dimensional Lie algebra $\mathfrak{g}$. Let

$$\begin{aligned} n_-(t_1, \dots, t_k) &:= e^{t_1 Y_1} e^{t_2 Y_2} \dots e^{t_k Y_k}, \\ a(t_{k+1}, \dots, t_{k+r}) &:= e^{t_{k+1} H_1} \dots e^{t_{k+r} H_r}, \text{ and} \\ n_+(t_{k+r+1}, \dots, t_d) &:= e^{t_{k+r+1} X_1} \dots e^{t_d X_k}. \end{aligned} \quad (\text{A.8})$$

⁶See, for example, the note <http://math.mit.edu/~stevenj/bump-saddle.pdf> (which uses a different normalization of Fourier transform).

Then for $c > 0$ sufficiently small the image of the set $\{|t_i| \leq c | 1 \leq i \leq d\}$ under the map

$$(t_1, t_2, \dots, t_d) \mapsto n_-(t_1, \dots, t_k) a(t_{k+1}, \dots, t_{k+r}) n_+(t_{k+r+1}, \dots, t_d) \quad (\text{A.9})$$

diffeomorphically sweeps out a compact neighborhood of the identity in G . Define a smooth function $\psi = \psi_c$ supported in this neighborhood by the formula

$$\psi_c(n_- a n_+) = \prod_{i=1}^d \sigma(c^{-1} t_i), \quad (\text{A.10})$$

where $n_- a n_+$ represents the righthand side of (A.9).

Let δ denote the right translation operator on functions on G . This action extends to $\mathfrak{g}$ as the left-invariant differential operators

$$(\delta(X)f)(g) = \left. \frac{d}{dt} \right|_{t=0} f(g e^{tX}), \quad X \in \mathfrak{g}, \quad (\text{A.11})$$

and then to the universal enveloping algebra $\mathfrak{U}$ of $\mathfrak{g}$ by composition.

Proposition. For any degree N element $X \in \mathfrak{U}$

$$\|\delta(X)\psi_c\|_1 \ll (c'N)^{c'N}, \quad (\text{A.12})$$

where $c' > 0$ is a constant that depends on c and G .

Proof. Let us first consider the case where $X \in \mathfrak{g}$ and the righthand side of (A.10) has the more general form $\prod_{i \leq d} \sigma_i(c^{-1} t_i)$, where $\sigma_1, \dots, \sigma_d \in C_c^\infty(\mathbb{R})$. Commuting the derivative in X across the n_+ factor

$$\begin{aligned} \delta(X)\psi_c(n_- a n_+) &= \left. \frac{d}{dt} \right|_{t=0} \psi_c(n_- a n_+ e^{tX}) \\ &= \left. \frac{d}{dt} \right|_{t=0} \psi_c(n_- a e^{t \text{Ad}(n_+)X} n_+). \end{aligned} \quad (\text{A.13})$$

Since $\psi_c(g)$ is supported on elements g which factorize as $g = n_- a n_+$, in which the support of each of the three factors n_- , a , and n_+ is constrained to a fixed compact set, we observe that $\text{Ad}(n_+)X$ can be written as a linear combination of elements of $\mathcal{B}$ with bounded coefficients. Furthermore, since $a \in A$ normalizes n_- the same compactness argument allows us to write

$$\delta(X)\psi_c(n_- a n_+) = \left. \frac{d}{dt} \right|_{t=0} \psi_c(n_- e^{tY'} a e^{tH'} e^{tX'} n_+), \quad (\text{A.14})$$

where $Y' \in \mathfrak{n}_-$, $H' \in \mathfrak{a}$, and $X' \in \mathfrak{n}_+$ depend on X , and range over bounded subsets as X varies over the basis $\mathcal{B}$. In particular, the coefficients of Y' , H' , or X' when expanded in the basis $\mathcal{B}$ are bounded by a constant M (depending only on c and G) as X ranges among the vectors $\mathcal{B}$. Thus for $X \in \mathcal{B}$

$$\begin{aligned} |\delta(X)\psi_c(n_- a n_+)| &\leq \\ M c^{-1} \sum_{i=1}^d &|\sigma_1(c^{-1} t_1) \cdots \sigma_{i-1}(c^{-1} t_{i-1}) \sigma'_i(c^{-1} t_i) \sigma_{i+1}(c^{-1} t_{i+1}) \cdots \sigma_d(c^{-1} t_d)|. \end{aligned} \quad (\text{A.15})$$

Similarly, when X is the tensor product of N basis vectors we have the estimate

$$|\delta(X)\psi_c(n_- a n_+)| \leq M^N c^{-N} \sum_{\substack{(j_1, \dots, j_d) \in \mathbb{Z}_{\geq 0}^d \\ j_1 + \dots + j_d = N}} \left| \sigma_1^{(j_1)}(c^{-1} t_1) \sigma_2^{(j_2)}(c^{-1} t_2) \cdots \sigma_d^{(j_d)}(c^{-1} t_d) \right|, \quad (\text{A.16})$$

as can be seen by iteration.

In the (compact) range of support of ψ , the Haar measure dg in the coordinates (A.9) is bounded by a multiple of $dt_1 \cdots dt_d$ and so by the lemma

$$\begin{aligned}
\|\delta(X)\psi_c\|_1 &\ll M^N c^{-N+d} \sum_{\substack{(j_1, \dots, j_d) \in \mathbb{Z}_{\geq 0}^d \\ j_1 + \dots + j_d = N}} \|\sigma_1^{(j_1)}\|_1 \cdots \|\sigma_d^{(j_d)}\|_1 \\
&\ll M^N c^{-N+d} \sum_{\substack{(j_1, \dots, j_d) \in \mathbb{Z}_{\geq 0}^d \\ j_1 + \dots + j_d = N}} (2j_1)^{2j_1} \cdots (2j_d)^{2j_d} \\
&\leq M^N c^{-N+d} \sum_{\substack{(j_1, \dots, j_d) \in \mathbb{Z}_{\geq 0}^d \\ j_1 + \dots + j_d = N}} (2N)^{2j_1} \cdots (2N)^{2j_d}.
\end{aligned} \tag{A.17}$$

The sum above is equal to the multinomial expansion of $\overbrace{(4N^2 + \cdots + 4N^2)}^{N \text{ times}} = (4N^3)^N$, and hence the lemma follows. $\square$

A.2.3. Next, we generalize the above estimate to the K -twists of the function ψ defined by

$$f_c(g) := \int_K \psi_c(k^{-1}gk) dk. \tag{A.18}$$

For any fixed $X \in \mathfrak{g}$ the elements $\text{Ad}(k^{-1})X$ are bounded as k ranges over the maximal compact subgroup K . In particular if $X \in \mathcal{B}$ then

$$(\delta(X)f_c)(g) = \left. \frac{d}{dt} \right|_{t=0} \int_K \psi_c(k^{-1}ge^{tX}k) dk \ll \sum_{X' \in \mathcal{B}} \int_K |(\delta(X')\psi_c)(k^{-1}gk)| dk, \tag{A.19}$$

with an implied constant that depends only on $K \subset G$.

Proposition. *There exists a constant $c' > 0$ which depends only on c and G such that*

$$\|\delta(X)f_c\|_1 \ll \sum_{X' \in \mathcal{B}} \|\delta(X')\psi_c\|_1 \ll (c'N)^{c'N} \tag{A.20}$$

for all $X \in \mathcal{B}$.

Proof. The first inequality follows from (A.20) using the bi-invariance of the Haar measure, and the second from proposition A.2.2. $\square$

A.2.4. A result of Harish-Chandra [18] (see also [3, p. 22]) asserts that for any K -finite automorphic form ϕ , there exists a function $f \in C_c^\infty(G)$ such that ϕ is in fact *equal* to the convolution

$$(\phi \star f)(g) := \int_G \phi(gh^{-1}) f(h) dh. \tag{A.21}$$

The proof furthermore shows that f satisfies the condition $f(k^{-1}gk) = f(g)$ for all $k \in K$, and can be taken to be a linear combination of functions in an approximate identity sequence. For ψ_c of the form (A.10), the K -twists defined in (A.18) are approximate identity sequences. This allows us to conclude the following

Proposition. *For any K -finite automorphic form ϕ there exists a finite number of positive real numbers $c_i > 0$ and complex numbers a_i such that*

$$\phi = \sum_i a_i (\phi \star f_{c_i}). \quad (\text{A.22})$$

For completeness we give the proof, which is completely based on ideas of Borel and Harish-Chandra as in *op. cit.*

Proof. As $c \rightarrow 0$ the functions f_c in (A.18) are smooth, nonnegative, and have support shrinking to zero. Thus for any $\phi_1 \in C_c^\infty(G)$ the convolutions $m_c^{-1}(\phi_1 \star f_c)$ converge to ϕ_1 uniformly on compacta, where $m_c = \|f_c\|_1$. At the same time, changing variables in (A.18) demonstrates that $f_c(k^{-1}gk) = f_c(g)$, and so $\phi \star f_c$ lies in $V :=$ the span of all K -translates of ϕ , a finite dimensional space.⁷ The linear span W of the convolution operators $\{\star f_c | c > 0\}$ is a subspace of $\text{Aut}(V)$ that is closed under any of its equivalent topologies, being that both are finite dimensional. The particular sequence of operators $\{\star(m_c^{-1}f_c)\}$ converges to the identity operator on V under the topology of uniform convergence on compacta, and hence the identity operator is in W ; that is, the identity operator on W has the form $\star \sum_i a_i f_{c_i}$, which implies (A.22). $\square$

A.2.5. We shall now combine the results of the previous paragraphs. According to (A.21),

$$\delta(X)(\phi \star f) = \phi \star \delta(X)f \quad (\text{A.23})$$

for any $X \in \mathfrak{U}$, as can be seen by changing variables. Thus the boundedness of cusp forms combined with estimate (A.20) shows that

$$\|\delta(X)(\phi \star f_c)\|_\infty \leq \|\phi\|_\infty \|\delta(X)f_c\|_1 \leq (c'N)^{c'N}, \quad (\text{A.24})$$

where N is the degree of X and $c' > 0$ is a constant depending on c and G . Proposition A.2.4 shows that ϕ is a finite linear combination of convolutions $\phi \star f_{c_i}$, and so we conclude

Proposition. *Let ϕ be a K -finite cusp form on G . There exists a positive constant $C > 0$ depending only on ϕ such that for every $X \in \mathfrak{U}$ of degree N*

$$\|\delta(X)\phi\|_\infty \leq (CN)^{CN}. \quad (\text{A.25})$$

A.3 Rapid decay with uniform constants

In this section we use proposition A.2.5 to give a rapid decay estimate with uniform constants. The methods here are fairly standard (see [27, §2.10], for example), though it is unfortunately necessary to reproduce them in order to demonstrate how the constant factors in (A.25) propagate.

A.3.1. Recall the conventions of §2.2 and §2.3, where we fixed an Iwasawa decomposition $G = UAK$ and a projection $\text{Iw}_A : G \rightarrow A$. Furthermore, in (2.7) we defined a character $A \rightarrow \mathbb{C}^*$ which sends $a \mapsto a^\mu$ for each linear functional $\mu : \mathfrak{h} = \text{Lie}(A) \rightarrow \mathbb{C}$.

⁷This is the only point in this appendix where the K -finiteness property is used.

Theorem. Let ϕ be a K -finite cusp form on $\Gamma \backslash G$. Fix a Siegel set $\mathfrak{S}_t$ for G . Then there exists a constant $C > 0$ which depends only on ϕ and the parameter t of the Siegel set $\mathfrak{S}_t$ such that

$$\phi(g) \leq (CN)^{CN} \text{Iw}_A(g)^{-N\rho} \quad \text{for all } g \in \mathfrak{S}_t \text{ and } N \geq 1. \quad (\text{A.26})$$

The remainder of this section will be devoted to the proof of this theorem.

A.3.2. Let $\alpha \in \Pi$ be a simple root and P the corresponding standard maximal parabolic subgroup, with unipotent radical U_P and corresponding Lie algebra $\mathfrak{n}_P$. Let $\Delta_{+,\alpha}$ denote the set of positive roots

$$\{\gamma \in \Delta_+ \mid \gamma = \sum_{\beta \in \Pi} m(\beta)\beta \text{ with } m(\alpha) \neq 0\}, \quad (\text{A.27})$$

which are precisely the roots γ whose Chevalley basis root vector X_γ lies in $\mathfrak{n}_P := \text{Lie}(U_P)$. Order the roots of $\Delta_{+,\alpha}$ according to their heights⁸:

$$\beta_1 > \beta_2 > \cdots > \beta_p.$$

For $i = 1, \dots, p$ let $X_i = X_{\beta_i}$ be the Chevalley basis root vector corresponding to the root β_i . Corresponding to the set $\{\beta_1, \dots, \beta_i\}$ for $0 \leq i \leq p$, we let $\mathfrak{n}_i \subset \mathfrak{n}_P$ and $U_i \subset U_P$ denote the corresponding nilpotent subalgebra and normal unipotent subgroups of $\mathfrak{n}_P$ and U_P , respectively; that is, $\mathfrak{n}_i$ is the real span of $\{X_1, \dots, X_i\}$ and U_i is its exponential. We then have the following ascending chains of Lie algebras and unipotent groups:

$$\{0\} = \mathfrak{n}_0 \subset \mathfrak{n}_1 \subset \cdots \subset \mathfrak{n}_p = \mathfrak{n}_P \quad (\text{A.28})$$

$$\{0\} = U_0 \subset U_1 \subset \cdots \subset U_p = U_P. \quad (\text{A.29})$$

For $1 \leq i \leq p$ the one parameter subgroups $\chi_{\beta_i}(s) = e^{sX_i}$ induce isomorphisms $U_{i-1} \backslash U_i \cong \mathbb{R}$; letting $\Gamma_\infty = \Gamma \cap U$, $\Gamma_i = \Gamma \cap U_i$, and rescaling the X_i if necessary, they furthermore induce isomorphisms

$$\chi_{\beta_i} : \mathbb{Z} \backslash \mathbb{R} \cong (U_{i-1} \Gamma_i) \backslash U_i \quad (\text{A.30})$$

that will be used to parametrize Fourier series below. Since replacing X_i by a positive multiple will make no difference in the rest of this appendix, we shall assume this rescaling has been performed.

A.3.3. For any function $\Psi \in L^1(\Gamma \backslash G)$ we can define projections

$$\Psi_i(g) := \int_{\Gamma_i \backslash U_i} \Psi(u_i g) du_i, \quad 0 \leq i \leq p, \quad (\text{A.31})$$

where du_i is the Haar measure on U_i normalized to give the quotient $\Gamma_i \backslash U_i$ measure one. These Haar measures can be written as products of Haar measures along the one-parameter subgroups given by the characters χ_{β_j} for $j \leq i$. Using (A.30) we can write

$$\Psi_i(g) = \int_{(U_{i-1} \Gamma_i) \backslash U_i} \Psi_{i-1}(ug) du = \int_{\mathbb{Z} \backslash \mathbb{R}} \Psi_{i-1}(\chi_{\beta_i}(s)g) ds$$

⁸One can choose any ordering $>$ on Δ with the property that $\beta, \gamma, \beta + \gamma \in \Delta \implies \beta + \gamma > \beta$ and $\beta + \gamma > \gamma$.

and hence

$$\begin{aligned}\Psi_i(g) - \Psi_{i-1}(g) &= \int_0^1 [\Psi_{i-1}(\chi_{\beta_i}(s)g) - \Psi_{i-1}(\chi_{\beta_i}(0)g)] ds \\ &= \int_0^1 \int_0^s \left(\frac{d}{dr} \Psi_{i-1}(\chi_{\beta_i}(r)g) \right) dr ds.\end{aligned}\tag{A.32}$$

Thus

$$|\Psi_i(g) - \Psi_{i-1}(g)| \leq \max_{0 \leq r \leq 1} \left| \frac{d}{dr} \Psi_{i-1}(\chi_{\beta_i}(r)g) \right| \tag{A.33}$$

is bounded in terms of derivatives of Ψ_{i-1} under *left* translation by the one-parameter subgroup generated by X_i .

We next describe these left derivatives in terms of the right action (A.11):

$$\frac{d}{dr} \Psi_{i-1}(\chi_{\beta_i}(r)g) = (\delta((\text{Ad } g^{-1})X_i) \Psi_{i-1})(\chi_{\beta_i}(r)g), \tag{A.34}$$

which can be seen by passing the Lie algebra derivative across the argument $\chi_{\beta_i}(r)g$ of Ψ_{i-1} . Suppose now that g has a factorization $g = uak$, in which u and k range over fixed compact sets; this description in particular applies to elements of Siegel sets. Because of our assumption on the ordering $>$, $\text{Ad}(u^{-1})X_{\beta_i}$ can be written as a linear combination of the root vectors $\{X_{\beta_1}, \dots, X_{\beta_j}\}$ with bounded coefficients; the precise bound depends on the size of the compact set u ranges over, and in particular depends only on Γ in the circumstance that $g \in \mathfrak{S}_t$. For each $a \in A$, $\text{Ad}(a^{-1})X_{\beta_j} = a^{-\beta_j}X_{\beta_j}$; because of (A.27) and the definition of $\mathfrak{S}_t$, the factor $a^{-\beta_j} \ll a^{-\alpha}$ with an implied constant depending only on t and G . Finally, $\text{Ad}(k^{-1})$ maps any fixed element of $\mathfrak{g}$ to a bounded set. Putting together these statements about $\text{Ad}(u^{-1})$, $\text{Ad}(a^{-1})$, and $\text{Ad}(k^{-1})$, we see that $\text{Ad}(g^{-1})X_i = \text{Ad}(k^{-1})\text{Ad}(a^{-1})\text{Ad}(u^{-1})X_i$ is a linear combination of elements of (A.7) with coefficients bounded by $a^{-\alpha}$ times a constant that depends only on the Siegel set $\mathfrak{S}_t$. Combining (A.33), (A.34), the sup-norm inequality $\|\delta(X)\Psi_i\|_\infty \leq \|\delta(X)\Psi\|_\infty$, and this expansion of $\text{Ad}(g^{-1})X_i$, we see that

$$|\Psi_{i-1}(g) - \Psi_i(g)| \ll a^{-\alpha} \max_{X \in \mathcal{B}} \|\delta(X)\Psi_i\|_\infty \leq a^{-\alpha} \max_{X \in \mathcal{B}} \|\delta(X)\Psi\|_\infty, \tag{A.35}$$

with an implied constant that depends only on $\mathfrak{S}_t$.

Specialize Ψ to be the cusp form ϕ , so that $\Psi_p = \phi_p = 0$. By applying the estimate (A.35) to the telescoping sum

$$\phi(g) = \sum_{i=1}^p (\phi_{i-1}(g) - \phi_i(g)), \tag{A.36}$$

we conclude that $\phi(g)$ is bounded by the product of $a^{-\alpha}$, the sup norms $\|\delta(X)\phi\|_\infty$ for any X in (A.7), and a constant that depends only on $\mathfrak{S}_t$. Theorem A.3.1 then follows from iterating this procedure for various simple roots α and applying the estimate in proposition A.2.5.

Recall that Theorem 4.6 is deduced from Theorem A.3.1 immediately after its statement on page 26.

A.4 Exponential decay

Lemma. *Let $F : \mathbb{R}_{>0} \rightarrow \mathbb{C}$ be a bounded function for which there exists constants $c > 0$ and $C > 0$ such that*

$$F(y) \leq cy^{-N} (CN)^{CN} \text{ for all } N \geq 1. \tag{A.37}$$

Then there exists constants $d > 0$ and $D > 0$ such that

$$F(y) \leq D \exp(-y^d). \quad (\text{A.38})$$

The converse is of course elementary. Theorem A.3.1 implies Theorem A.1 as a consequence of this lemma, and also vice-versa (though with different constants).

Proof. By adjusting the constants d and D it suffices to show the weaker bound $F(y) \leq Dye^{-y^d}$ for $y > 1$. If $y > 1$ the assumption (A.37) implies

$$y^{dn-1} F(y) \leq y^{[dn]} F(y) \leq c(C[dn])^{C[dn]} \leq c(Cdn)^{Cdn} \quad (\text{A.39})$$

for any $d > 0$ and $n \in \mathbb{Z}_{>0}$. For d sufficiently small the righthand side is less than a constant times $(n!)/n^2$, so summing over n gives the boundedness of

$$y^{-1} \exp(y^d) F(y) = \sum_{n=0}^{\infty} \frac{1}{n!} y^{dn-1} F(y) \quad (\text{A.40})$$

in the range $y > 1$, as was to be shown. $\square$

A.5 Some rapidly, but not exponentially, decaying smooth cusp forms

In this section we give an example of a non- K -finite yet smooth cusp form which does not satisfy the exponential-type decay statement present in Theorem A.1. It is a theorem of [26] that all smooth cusp forms have rapid decay, but the proof does not give the uniformity of constants present in the stronger statement (A.26) from Theorem A.3.1. The example here shows that not only is this uniformity impossible, but furthermore that its failure is very common: this is because we show that the decay is governed by an arbitrarily chosen Schwartz function (relatively few of which decay faster than an exponential).

We give an example for the group $SL(2, \mathbb{R})$, though the phenomenon is certainly much more general. Let π be any cuspidal automorphic representation for $SL(2, \mathbb{Z}) \backslash SL(2, \mathbb{R})$. We will demonstrate it has non- K -finite smooth vectors which violate (A.1). To do this it is convenient to use the notion of *automorphic distribution* as in [25]. The representation π has an automorphic distribution τ which embeds smooth vectors v for some principal series representation⁹ to automorphic forms ϕ :

$$\phi(n_x a_y) = \langle \tau, \pi(n_x a_y) v \rangle, \text{ where } n_x = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \text{ and } a_y = \begin{pmatrix} y^{1/2} & 0 \\ 0 & y^{-1/2} \end{pmatrix}. \quad (\text{A.41})$$

In the line model, the Schwartz space is an embedded subspace of smooth vectors. If v corresponds to a Schwartz function $f(u)$, then $\pi(n_x)v$ corresponds to $f(u-x)$ and $\pi(a_y)v$ corresponds to $f(\frac{u}{y})|y|^\mu \text{sgn}(a)^\delta$, where $\mu \in \mathbb{C}$ and $\delta \in \mathbb{Z}/2\mathbb{Z}$ depend on the archimedean type of the representation π . The distribution τ is periodic, hence tempered (meaning that it can be integrated against arbitrary Schwartz functions). Its restriction to the real line has a Fourier series,

$$\tau(u) = \sum_{n \neq 0} c_n e^{2\pi i n u}, \quad (\text{A.42})$$

⁹This is the case even if the representation is not itself principal series, e.g., discrete series.

with coefficients that grow at most polynomially in n (see [25]). Thus

$$\begin{aligned}
\phi(n_x a_y) &= |y|^\mu \operatorname{sgn}(y)^\delta \int_{\mathbb{R}} \sum_{n \neq 0} c_n e^{2\pi i n u} f\left(\frac{u-x}{y}\right) du \\
&= |y|^{\mu+1} \operatorname{sgn}(y)^\delta \sum_{n \neq 0} c_n e^{2\pi i n x} \int_{\mathbb{R}} e^{2\pi i n y u} f(u) du \\
&= |y|^{\mu+1} \operatorname{sgn}(y)^\delta \sum_{n \neq 0} c_n e^{2\pi i n x} \widehat{f}(-ny).
\end{aligned} \tag{A.43}$$

The estimate (A.1) implies exponential decay (of the same caliber) for each Fourier coefficient of ϕ , in particular the first coefficient. In particular, it implies $\widehat{f}(-y)$ decays faster than the exponential of some fractional power of y . However, it is an arbitrary Schwartz function, and not all Schwartz functions have that decay property, e.g. $y \mapsto e^{-(\ln(y^2+1))^2}$. We conclude that the exponential decay estimate (A.1) does not hold for all smooth cusp forms.

A.6 Exponential decay estimates in string theory

A number of interesting quantities in string theory are, by their very definition, automorphic functions. This is because split exceptional real groups in the E_n series arise as duality groups in toroidal compactifications in Cremmer-Julia supergravity, and as a consequence the coupling constants of the theory lie in a symmetric space G/K . In type IIB string theory, S -, T -, and U -dualities force invariance under a large discrete subgroup $\Gamma \subset G$ [20]. The purpose of this section is to discuss the exact exponent d in the estimate (A.1), and its string theory interpretation in various situations. Much of the discussion here is an outgrowth of the second-named author's joint work with Michael Green and Pierre Vanhove [16], who we thank for their insightful comments on the questions posed here. More information about these questions can be found in [1, 16, 17, 28, 29] and in related references therein.

A.6.1. Decay estimates for Fourier coefficients

In various particular problems supersymmetry provides extra information such as differential equations (e.g., for the Laplace-Beltrami operator), suggesting that the automorphic functions are automorphic forms in these situations. In addition, at times supersymmetry predicts the asymptotic behaviour of various Fourier coefficients of these automorphic forms.

Although the exponential decay estimate (A.1) is stated for cuspidal automorphic forms ϕ , it of course applies to any Fourier coefficient defined by integration of a character over a unipotent subgroup $U' \subset U$:

$$\phi_\chi(g) := \int_{(\Gamma \cap U') \backslash U'} \phi(ug) \chi(u)^{-1} du, \tag{A.44}$$

where du is the Haar measure normalized to give the quotient $(\Gamma \cap U') \backslash U'$ volume 1 and χ is a character of U which is trivial on $\Gamma \cap U'$. Indeed, since the integration is over a compact quotient ϕ_χ also obeys the estimate (A.1). Thus Theorem A.1 gives the exponential decay of these Fourier coefficients. When ϕ is not cuspidal similar decay is expected for nontrivial characters χ , since an automorphic form minus its constant terms has rapid decay (this was essentially rederived in the analysis of (A.36) above).

A.6.2. Some precise questions on the exponential decay for maximal parabolics

Let α denote a positive simple root and $P = P_\alpha \supset B$ denote its associated standard maximal parabolic subgroup of G , which has the factorization $P_\alpha = A_\alpha M_\alpha U_\alpha$ in terms of its unipotent radical U_α and the center A_α of its reductive part $A_\alpha M_\alpha$. For $a \in A_\alpha$ recall that $a^\alpha \in \mathbb{R}$ is defined by the formula $\text{Ad}(a)X_\alpha = a^\alpha X_\alpha$, where X_α is a root vector for the simple root α . We shall use this coordinate to describe limits along the group A_α , which often has an interpretation in string theory, e.g., of a compactifying radius or string coupling constant. We do not assume that ϕ is a cusp form but we do assume that χ is nontrivial in order to exclude constant terms.

Question A: Is it always the case that there exists constants $c_1 = c_1(g, \phi, \chi)$, $c_2 = c_2(g, \phi, \chi)$, and $d = d(\phi, \chi)$ such that

$$\phi_\chi(ag) \simeq c_1(g, \phi, \chi) \exp\left(-c_2(g, \phi, \chi)(a^\alpha)^{d(\phi, \chi)}\right) \quad (\text{A.45})$$

for g fixed, as $a^\alpha \rightarrow \infty$? Here the symbol $\simeq$ is used to indicate asymptotic equality up to polynomials in the a^α . (Such factors do arise, e.g., in [16, (H.31)]).

Question B: Assuming an affirmative answer to Question A, does $d(\phi, \chi)$ depend nontrivially on χ , or is instead always 1 (as it is in all currently known cases)?

Let us now consider the lower central series

$$U_\alpha = U^{(1)} \supset U^{(2)} \supset U^{(3)} \supset \dots \supset U^{(\ell)} = \{e\}, \quad (\text{A.46})$$

where $U^{(k+1)} = [U^{(k)}, U_\alpha]$.

Question C: Fix $1 \leq k \leq \ell$ and a nontrivial character χ of U_k which is trivial on $\Gamma \cap U^{(k)}$. Assuming an affirmative answer to Question A, is $d(\phi, \chi) = k$?

For example, when U_α is a Heisenberg group Question C asks whether or not nontrivial Fourier coefficients along the one-dimensional center of U_α decay significantly faster than nontrivial Fourier coefficients on the full group U_α do. We present some motivation for this question below.

A.6.3. Examples from string theory

We shall now describe the origin of question C in string theory. Let G denote the split real form of the simply connected Chevalley group of type E_n , where $n = 6, 7$, or 8 . Let α denote the last positive simple root of G in the usual Bourbaki numbering. Thus M_α is a simply connected Chevalley group of type $\text{Spin}(5, 5)$, E_6 , or E_7 in these three cases.

For particular Eisenstein series (which are not cuspidal, but for which our arguments nevertheless give the same bounds on Fourier coefficients) the nonvanishing Fourier coefficients are expected to obey precise asymptotics as a^α goes to infinity. This is because a^α corresponds to the decompactifying radius in the toroidal compactification. When such a radius tends to infinity, the compactified theory resembles that of a higher dimensional theory compactified on a lower-dimensional torus, and has duality group M_α . Precise statements often give exponential decay in terms of an effective action, which in our context implies that $d(\phi, \chi) = 1$ when U_α is abelian and α comes from one of the three terminal nodes of G 's Dynkin diagram. This motivates Question B and the $k = 1$ case of Question C.

The string theory interpretation is apparently different if U_α is nonabelian, for example a Heisenberg group. In this case there is an additional charge to consider, which is instead consistent with $d(\phi, \chi) = 2$ for the $k = 2$ case of Question C. Evidence for this is presented in [1, 28, 29].

It should be stressed that string theory does not provide any additional insight into the decay rate of general automorphic forms, other than pointing to subtleties previously unrecognized in particular examples.

References

- [1] L. Bao, A. Kleinschmidt, B. E. W. Nilsson, D. Persson, and B. Pioline, *Instanton Corrections to the Universal Hypermultiplet and Automorphic Forms on $SU(2,1)$* , Commun. Num. Theor. Phys. **4** (2010), 187–266. arXiv:0909.4299 [hep-th].
- [2] Armand Borel, *Introduction aux groupes arithmétiques*, Publications de l’Institut de Mathématique de l’Université de Strasbourg, XV. Actualités Scientifiques et Industrielles, No. 1341, Hermann, Paris, 1969 (French). MR0244260 (39 #5577)
- [3] ———, *Automorphic forms on $SL_2(\mathbf{R})$* , Cambridge Tracts in Mathematics, vol. 130, Cambridge University Press, Cambridge, 1997. MR1482800 (98j:11028)
- [4] A. Braverman and D. Kazhdan, *Representations of affine Kac-Moody groups over local and global fields: a survey of some recent results*. arXiv:1205.0870.
- [5] Howard Garland, *The arithmetic theory of loop algebras*, J. Algebra **53** (1978), no. 2, 480–551, DOI 10.1016/0021-8693(78)90294-6. MR502647 (80a:17012)
- [6] ———, *The arithmetic theory of loop groups*, Inst. Hautes Études Sci. Publ. Math. **52** (1980), 5–136. MR601519 (83a:20057)
- [7] ———, *The arithmetic theory of loop groups. II. The Hilbert-modular case*, J. Algebra **209** (1998), no. 2, 446–532, DOI 10.1006/jabr.1998.7529. MR1659899 (2000h:20088)
- [8] ———, *Certain Eisenstein series on loop groups: convergence and the constant term*, Algebraic groups and arithmetic, Tata Inst. Fund. Res., Mumbai, 2004, pp. 275–319. MR2094114 (2005i:11118)
- [9] ———, *Absolute convergence of Eisenstein series on loop groups*, Duke Math. J. **135** (2006), no. 2, 203–260, DOI 10.1215/S0012-7094-06-13521-4. MR2267283 (2008g:22032)
- [10] ———, *Eisenstein series on loop groups: Maass-Selberg relations. I*, Algebraic groups and homogeneous spaces, Tata Inst. Fund. Res. Stud. Math., Tata Inst. Fund. Res., Mumbai, 2007, pp. 275–300. MR2348907 (2008j:22029)
- [11] ———, *Eisenstein series on loop groups: Maass-Selberg relations. II*, Amer. J. Math. **129** (2007), no. 3, 723–784, DOI 10.1353/ajm.2007.0017. MR2325102 (2008j:22030)
- [12] ———, *Eisenstein series on loop groups: Maass-Selberg relations. III*, Amer. J. Math. **129** (2007), no. 5, 1277–1353, DOI 10.1353/ajm.2007.0032. MR2354321 (2008j:22031)
- [13] ———, *Eisenstein series on loop groups: Maass-Selberg relations. IV*, Lie algebras, vertex operator algebras and their applications, Contemp. Math., vol. 442, Amer. Math. Soc., Providence, RI, 2007, pp. 115–158, DOI 10.1090/conm/442/08522, (to appear in print). MR2372559 (2008j:22032)
- [14] ———, *On extending the Langlands-Shahidi method to arithmetic quotients of loop groups*, Representation theory and mathematical physics, Contemp. Math., vol. 557, Amer. Math. Soc., Providence, RI, 2011, pp. 151–167, DOI 10.1090/conm/557/11030, (to appear in print). MR2848924
- [15] Roger Godement, *Introduction à la théorie de Langlands*, Séminaire Bourbaki, Vol. 10, Soc. Math. France, Paris, 1995, pp. Exp. No. 321, 115–144 (French). MR1610464
- [16] Michael B. Green, Stephen D. Miller, and Pierre Vanhove, *Small representations, string instantons, and Fourier modes of Eisenstein series*. arxiv:1111.2983.
- [17] Michael B. Green, Stephen D. Miller, Jorge G. Russo, and Pierre Vanhove, *Eisenstein series for higher-rank groups and string theory amplitudes*, Commun. Number Theory Phys. **4** (2010), no. 3, 551–596. MR2771579

- [18] Harish-Chandra, *Automorphic forms on a semisimple Lie group*, Proc. Nat. Acad. Sci. U.S.A. **45** (1959), 570–573. MR0105461 (21 #4202)
- [19] ———, *Automorphic forms on semisimple Lie groups*, Notes by J. G. M. Mars. Lecture Notes in Mathematics, No. 62, Springer-Verlag, Berlin, 1968. MR0232893 (38 #1216)
- [20] C. M. Hull and P. K. Townsend, *Unity of Superstring Dualities*, Nucl. Phys. B **438** (1995), 109–137. arXiv:hep-th/9410167.
- [21] N. Iwahori and H. Matsumoto, *On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups*, Inst. Hautes Études Sci. Publ. Math. **25** (1965), 5–48. MR0185016 (32 #2486)
- [22] Victor G. Kac, *Infinite-dimensional Lie algebras*, 3rd ed., Cambridge University Press, Cambridge, 1990. MR1104219 (92k:17038)
- [23] Bernhard Krötz and Eric Opdam, *Analysis on the crown domain*, Geom. Funct. Anal. **18** (2008), no. 4, 1326–1421, DOI 10.1007/s00039-008-0684-5. MR2465692 (2010a:22011)
- [24] Robert P. Langlands, *Euler products*, Yale University Press, New Haven, Conn., 1971. A James K. Whittemore Lecture in Mathematics given at Yale University, 1967; Yale Mathematical Monographs, 1. MR0419366 (54 #7387)
- [25] Stephen D. Miller and Wilfried Schmid, *Summation formulas, from Poisson and Voronoi to the present*, Non-commutative harmonic analysis, Progr. Math., vol. 220, Birkhäuser Boston, Boston, MA, 2004, pp. 419–440. MR2036579 (2004m:11075)
- [26] ———, *On the rapid decay of cuspidal automorphic forms*, Adv. Math. **231** (2012), no. 2, 940–964, DOI 10.1016/j.aim.2012.06.001. MR2955198
- [27] C. Mœglin and J.-L. Waldspurger, *Spectral decomposition and Eisenstein series*, Cambridge Tracts in Mathematics, vol. 113, Cambridge University Press, Cambridge, 1995. Une paraphrase de l’Écriture [A paraphrase of Scripture]. MR1361168 (97d:11083)
- [28] D. Persson, *Automorphic Instanton Partition Functions on Calabi-Yau Threefolds*. arXiv:1103.1014 [hep-th].
- [29] P. Pioline and D. Persson, *The Automorphic NS5-brane*, Commun. Num. Theor. Phys. **3** (2009), 697–754. arXiv:0902.3274 [hep-th].
- [30] M. S. Raghunathan, *On the first cohomology of discrete subgroups of semisimple Lie groups*, Amer. J. Math. **87** (1965), 103–139. MR0173730 (30 #3940)
- [31] Freydoon Shahidi, *Eisenstein series and automorphic L -functions*, American Mathematical Society Colloquium Publications, vol. 58, American Mathematical Society, Providence, RI, 2010. MR2683009 (2012d:11119)
- [32] F. Sunyer Balaguer, *Sobre los momentos de las funciones holomorfas y acotadas en un angulo*, Revista Matemática Hispano-Americana **13** (1953), no. 4, 241–246.

e-mail addresses: garland-howard@yale.edu, miller@math.rutgers.edu,
patnaik@ualberta.ca